

MIXING OPERATORS AND SMALL SUBSETS OF THE CIRCLE

FRÉDÉRIC BAYART AND ÉTIENNE MATHERON

ABSTRACT. We provide complete characterizations, on Banach spaces with cotype 2, of those linear operators which happen to be weakly mixing or strongly mixing transformations with respect to some nondegenerate Gaussian measure. These characterizations involve two families of small subsets of the circle: the countable sets, and the so-called *sets of uniqueness* for Fourier-Stieltjes series. The most interesting part, i.e. the sufficient conditions for weak and strong mixing, is valid on an arbitrary (complex, separable) Fréchet space.

1. INTRODUCTION

1.1. Ergodic measures for linear operators. A basic problem in topological dynamics is to determine whether a given continuous map $T : X \rightarrow X$ acting on a topological space X admits an ergodic probability measure. One may also ask for stronger ergodicity properties such as weak mixing or strong mixing, and put additional constraints on the measure μ , for example that μ should have no discrete part, or that it should belong to some natural class of measures related to the structure of the underlying space X . Especially significant is the requirement that μ should have *full support* (i.e. $\mu(V) > 0$ for every open set $V \neq \emptyset$) since in this case any ergodicity property implies its topological counterpart. There is, of course, a huge literature on these matters since the classical work of Oxtoby and Ulam ([48], [49]).

In recent years, the above problem has received a lot of attention in the specific setting of *linear* dynamics, i.e. when the transformation T is a continuous linear operator acting on a topological vector space X ([27], [8], [7], [14], [30]). This is rather natural since the main objects of study in the area are the so-called *hypercyclic* operators, i.e. the operators having dense orbits. When the space X is second-countable, it is very easy to see that if a continuous map $T : X \rightarrow X$ happens to be ergodic with respect to some Borel probability measure μ with full support, then almost every $x \in X$ (relative to μ) has a dense T -orbit. Hence, to find an ergodic measure with full support is an efficient way of showing that a given operator is hypercyclic, which comes as a measure-theoretic counterpart to the more traditional Baire category approach.

2000 *Mathematics Subject Classification.* 47A16, 37A25.

Key words and phrases. Hypercyclic operators, Gaussian measures, mixing, unimodular eigenvalues, countable sets, sets of extended uniqueness, cotype 2.

Throughout the paper, we shall restrict ourselves to the best understood infinite-dimensional measures, the so-called *Gaussian* measures. Moreover, the underlying topological vector space X will always be a complex separable Fréchet space. (The reason for considering *complex* spaces only will become clear in the next few lines). In this setting, a Borel probability measure on X is Gaussian if and only if it is the distribution of an almost surely convergent random series of the form $\xi = \sum_0^\infty g_n x_n$, where $(x_n) \subset X$ and (g_n) is a sequence of independent, standard complex Gaussian variables. Given any property (P) relative to measure-preserving transformations, we shall say that an operator $T \in \mathfrak{L}(X)$ has property (P) *in the Gaussian sense* if there exists some Gaussian probability measure μ on X with full support with respect to which T has (P).

The problem of determining which operators are ergodic in the Gaussian sense was investigated by E. Flytzanis ([27]), in a Hilbert space setting. The fundamental idea of [27] is that one should look at the \mathbb{T} -*eigenvectors* of the operator, i.e. the eigenvectors associated with eigenvalues of modulus 1: roughly speaking, ergodicity is equivalent to the existence of “sufficiently many \mathbb{T} -eigenvectors and eigenvalues”. This is of course to be compared with the now classical eigenvalue criterion for hypercyclicity found by G. Godefroy and J. Shapiro ([29]), which says in essence that an operator having enough eigenvalues inside and outside the unit circle must be hypercyclic.

Building on Flytzanis’ ideas, the first named author and S. Grivaux came rather close to characterizing the weak and strong mixing properties for Banach space operators in terms of the \mathbb{T} -eigenvectors ([7], [8]). However, this was not quite the end of the story because the sufficient conditions for weak or strong mixing found in [7] and [8] depend on some geometrical property of the underlying Banach space, or on some “regularity” property of the \mathbb{T} -eigenvectors (see the comments just after Corollary 1.2).

In the present paper, our aim is to show that in fact, these assumptions can be completely removed. Thus, we intend to establish “optimal” sufficient conditions for weak and strong mixing in terms of the \mathbb{T} -eigenvectors which are valid on an arbitrary Fréchet space. These conditions turn out to be also necessary when the underlying space X is a Banach space with *cotype 2*, and hence we get complete characterizations of weak and strong mixing in this case.

1.2. Main results. Before stating our main results, we should recall the definitions of weak and strong mixing. A measure-preserving transformation $T : (X, \mathfrak{B}, \mu) \rightarrow (X, \mathfrak{B}, \mu)$ is *weakly mixing* (with respect to μ) if

$$\frac{1}{N} \sum_{n=0}^{N-1} |\mu(A \cap T^{-n}(B)) - \mu(A)\mu(B)| \xrightarrow{N \rightarrow \infty} 0$$

for any measurable sets $A, B \subset X$; and T is *strongly mixing* if

$$\mu(A \cap T^{-n}(B)) \xrightarrow{n \rightarrow \infty} \mu(A)\mu(B)$$

for any $A, B \in \mathfrak{B}$. (Ergodicity can be defined exactly as weak mixing, but removing the absolute value in the Cesàro mean).

According to the “spectral viewpoint” on ergodic theory, weakly mixing transformations are closely related to *continuous* measures on the circle \mathbb{T} (measures having no discrete part), and strongly mixing transformations are related to *Rajchman* measures (measures whose Fourier coefficients vanish at infinity). Without going into any detail at this point, we just recall that, by a classical result of Wiener (see e.g. [38]), continuous measures on \mathbb{T} are characterized by the behaviour of their Fourier coefficients: a measure σ is continuous if and only if

$$\frac{1}{N} \sum_{n=0}^{N-1} |\widehat{\sigma}(n)| \xrightarrow{N \rightarrow \infty} 0.$$

Wiener’s lemma is usually stated with symmetric Cesàro means, but this turns out to be equivalent. Likewise, by the so-called *Rajchman’s lemma*, a measure σ is Rajchman if and only if $\widehat{\sigma}(n) \rightarrow 0$ as $n \rightarrow +\infty$ (that is, a one-sided limit is enough).

Especially important for us will be the corresponding families of “small” sets of the circle; that is, the sets which negligible for every positive measure in the family under consideration (continuous measures, or Rajchman measures). Since this will have some importance later on, we note that negligibility with respect to some measure makes sense for arbitrary sets; in particular, negligible sets need not be Borel.

It follows from the classical Perfect Set Theorem that a Borel set $D \subset \mathbb{T}$ is small for continuous measures if and only if it is countable. As for Rajchman measures, the corresponding small sets are the so-called *sets of extended uniqueness* or *sets of uniqueness for Fourier-Stieltjes series*, which have been extensively studied since the beginning of the 20th century (see e.g. [35] or [40]). The family of all sets of extended uniqueness is usually denoted by \mathcal{U}_0 .

Our main results can now be summarized as follows.

Theorem 1.1. *Let X be a complex separable Fréchet space, and let $T \in \mathfrak{L}(X)$.*

- (1) *Assume that the \mathbb{T} -eigenvectors are perfectly spanning, in the following sense: for any countable set $D \subset \mathbb{T}$, the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in X . Then T is weakly mixing in the Gaussian sense.*
- (2) *Assume that the \mathbb{T} -eigenvectors are \mathcal{U}_0 -perfectly spanning, in the following sense: for any Borel set of extended uniqueness $D \subset \mathbb{T}$, the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in X . Then T is strongly mixing in the Gaussian sense.*
- (3) *In (1) and (2), the converse implications are true if X is a Banach space with cotype 2.*

Thus, we see in particular that if the \mathbb{T} -eigenvectors of T are perfectly spanning, then one can find a sequence $(x_n) \subset X$ spanning a dense subspace of X such that the Gaussian series $\xi = \sum g_n x_n$ is almost surely convergent, the distribution of ξ is T -invariant, and the random vector is almost surely hypercyclic for T .

Some remarks are in order regarding these results.

- Remarks.* (i) Theorem 1.1 says nothing about *ergodicity* in the Gaussian sense. This is because ergodicity in the Gaussian sense is in fact equivalent to weak mixing (see e.g. [28], or [8, Theorem 4.1]; see also Remark 1 after Lemma 3.4).
- (ii) The importance of the \mathbb{T} -eigenvectors is easy to explain. Indeed, it is almost trivial that if $T \in \mathfrak{L}(X)$ is an operator whose \mathbb{T} -eigenvectors span a dense subspace of X , then T admits an invariant Gaussian measure with full support: choose a sequence of \mathbb{T} -eigenvectors $(x_n)_{n \geq 0}$ (say $T(x_n) = \lambda_n x_n$) with dense linear span such that $\sum_0^\infty \|x_n\| < \infty$ for every continuous semi-norm $\|\cdot\|$ on X , and let μ be the distribution of the random variable $\xi = \sum_0^\infty g_n x_n$. That μ is T -invariant follows from the linearity of T and the rotational invariance of the Gaussian variables g_n ($\mu \circ T^{-1} \sim \sum_0^\infty g_n T(x_n) = \sum_0^\infty (\lambda_n g_n) x_n \sim \sum_0^\infty g_n x_n = \mu$). However, this particular measure μ cannot be ergodic ([30]).
- (iii) When X is a Hilbert space, (1) is stated in [27] (with some additional assumptions on the operator T) and a detailed proof is given in [7] (without these additional assumptions). The definition of “perfectly spanning” used in [7] is formally stronger than the above one, but the two notions are in fact equivalent ([30]).
- (iv) Recall that an operator $T \in \mathfrak{L}(X)$ is said to be *frequently hypercyclic* if there exists some vector $x \in X$ whose T -orbit visits every non-empty open set along a set of integers having positive lower density. (This notion was introduced in [7] and extensively studied since then; see e.g. the books [9] and [33] for more information). It follows from the pointwise ergodic theorem that ergodicity with respect to some probability measure with full support implies frequent hypercyclicity. In [30], it is shown that under the assumption of (1), the operator T is frequently hypercyclic. Actually, it follows from the main result of [30] and the ergodic decomposition theorem that under the perfect spanning assumption, the operator T admits an ergodic measure μ with full support. The proof of this result is rather complicated. It is not clear that it could be modified to get weak mixing with respect to this measure μ , and not clear either that it could produce a Gaussian measure in the end (which would yield weak mixing in the Gaussian sense). However, some of the ideas of [30] will be crucial for us. In particular, sub-section 3.3 owes a lot to [30].
- (v) In the weak mixing case, (3) is proved in [8, Theorem 4.1].
- (vi) It seems unnecessary to recall here the definition of cotype (see any book on Banach space theory, e.g. [2]). Suffices it to say that this is a geometrical property of the space, and that Hilbert space has cotype 2 as well as L^p spaces for $p \in [1, 2]$ (but *not* for $p > 2$).
- (vii) As shown in [8, Example 4.2], (3) does not hold on an arbitrary Banach space X . See section 7 (Kalisch operators) for more details.

For concrete applications, it is useful to formulate Theorem 1.1 in terms of \mathbb{T} -*eigenvector fields* for the operator T . A \mathbb{T} -eigenvector field for T is a map $E : \Lambda \rightarrow X$ defined on some set $\Lambda \subset \mathbb{T}$, such that

$$TE(\lambda) = \lambda E(\lambda)$$

for every $\lambda \in \Lambda$. (The terminology is not perfectly accurate: strictly speaking, $E(\lambda)$ is perhaps not a \mathbb{T} -eigenvector because it is allowed to be 0). Recall also that a closed set $\Lambda \subset \mathbb{T}$ is *perfect* if it has no isolated points or, equivalently, if $V \cap \Lambda$ is uncountable for any open set $V \subset \mathbb{T}$ such that $V \cap \Lambda \neq \emptyset$. Analogously, a closed set $\Lambda \subset \mathbb{T}$ is said to be \mathcal{U}_0 -*perfect* if $V \cap \Lambda$ is not a set of extended uniqueness for any open set V such that $V \cap \Lambda \neq \emptyset$. (For example, any nontrivial closed arc is \mathcal{U}_0 -perfect).

Corollary 1.2. *Let X be a separable complex Fréchet space, and let $T \in \mathfrak{L}(X)$. Assume that one has at hand a family of continuous \mathbb{T} -eigenvector fields $(E_i)_{i \in I}$ for T , where $E_i : \Lambda_i \rightarrow X$ is defined on some closed set $\Lambda_i \subset \mathbb{T}$, such that $\text{span}(\bigcup_{i \in I} E_i(\Lambda_i))$ is dense in X .*

- (i) *If each Λ_i is a perfect set, then T is weakly mixing in the Gaussian sense.*
- (ii) *If each Λ_i is \mathcal{U}_0 -perfect, then T is strongly mixing in the Gaussian sense.*

Proof. If $\Lambda \subset \mathbb{T}$ is a perfect set then $\Lambda \setminus D$ is dense in Λ for any countable set D , whereas if Λ is \mathcal{U}_0 -perfect then $\Lambda \setminus D$ is dense in Λ for any \mathcal{U}_0 -set D . Since the \mathbb{T} -eigenvector fields E_i are assumed to be continuous, it follows that the \mathbb{T} -eigenvectors of T are perfectly spanning in case (i), and \mathcal{U}_0 -perfectly spanning in case (ii). \square

Remark. The assumption in (i), i.e. the existence of \mathbb{T} -eigenvector fields with the required spanning property defined on perfect sets, is in fact equivalent to the perfect spanning property ([30]). Likewise, the assumption in (ii) is equivalent to the \mathcal{U}_0 -perfect spanning property (see Proposition 3.7).

1.3. Some non-technical comments. Several results similar to Corollary 1.2 are proved in [8] and in [9, Chapter 5], all of them being based on an interplay between the geometry of the (Banach) space X and the regularity of the \mathbb{T} -eigenvector fields E_i . For example, it is shown that if X has *type 2*, then continuity of the E_i is enough, whereas if the E_i are Lipschitz and defined on (nontrivial) closed arcs then no assumption on X is needed. “Intermediate” cases involve the type of the Banach space X and Hölder conditions on the E_i . What Corollary 1.2 says is that continuity of the E_i is *always* enough, regardless of the underlying space X .

A few more words are in order to explain in non-technical terms the main idea underlying the proof of this result.

To construct an ergodic or mixing Gaussian measure with full support $\mu \sim \sum g_n x_n$ for a given operator $T \in \mathfrak{L}(X)$, the “standard” procedure is to define the vectors x_n as Fourier coefficients of some suitable \mathbb{T} -eigenfield $E : \mathbb{T} \rightarrow X$ for T . The Fourier coefficients are relative to some orthonormal basis (e_n) of $L^2(\mathbb{T}, \sigma)$, where σ is a probability measure on \mathbb{T} . (In most concrete cases, σ is Lebesgue measure and (e_n) is the usual trigonometric basis). The main difficulty is to ensure at the same time that the Gaussian series $\sum g_n \widehat{E}(n)$ is almost surely convergent and that the $\widehat{E}(n)$ span a dense subspace of X .

Without any geometrical assumption on the Banach space X , the only way that comes to mind to ensure without fail the convergence of $\sum g_n \widehat{E}(n)$ is by requiring a strong “regularity” property of the vector field E , namely that $\sum \|\widehat{E}(n)\| < \infty$.

But if we stay within the very rigid framework of harmonic analysis *on the circle group* \mathbb{T} , there seems to be no hope to find such a regular \mathbb{T} -eigenfield E with the required spanning property.

On the other hand, it turns out that this can be done if we forget the circle group and replace it with a more “flexible” group, namely the *Cantor group* $G = \{0, 1\}^{\mathbb{N}}$. This is the basic idea of our proof in the weak mixing case. For the strong mixing property, things are technically harder, and more complicated “group-like” objects are needed; but the basic idea remains the same.

1.4. Organization of the paper. We have deliberately stressed the formal analogy between weak and strong mixing in the statement of Theorem 1.1. In view of this analogy, it should not come as a surprise that Theorem 1.1 can be deduced from some more general results dealing with abstract notions of “mixing”. (In order not to make this introduction exceedingly long, these results will be described in the next section). In particular, (1) and (2) are formal consequences of Theorem 2.6 below. However, even though the proof of Theorem 2.6 is “conceptually” simple, the technical details make it rather long. This would be exactly the same for the strong mixing case (i.e. part (2) of Theorem 1.1), but in the weak mixing case it is possible to give a technically much simpler and hence much shorter proof. For the sake of readability, it seems desirable to present this proof separately. But since there is no point in repeating identical arguments, we shall follow the abstract approach as long as this does not appear to be artificial.

The paper is organized as follows. In section 2, we present our abstract results. In section 3, we review some basic facts concerning Gaussian measures and we outline the strategy for proving the abstract results and hence Theorem 1.1. Apart from some details in the presentation and the level of generality, this follows the scheme described in [7], [8] and [9]. In section 4, we prove part (1) of Theorem 1.1 (the sufficient condition for weak mixing). The abstract results are proved in sections 5 and 6. Section 7 contains some concrete examples, including new ones (weighted shifts on $c_0(\mathbb{N})$, and Cowen-Douglas operators). Finally, miscellaneous remarks are collected in section 8. In particular, we briefly discuss the “continuous” analogues of our results, i.e. the case of 1-parameter semigroups (with an application to translation semigroups); we prove that operators satisfying the so-called Frequent Hypercyclicity Criterion are strongly mixing in the Gaussian sense; and we show that for a large class of strongly mixing weighted shifts, the set of hypercyclic vectors turns out to be rather small, namely Haar-null in the sense of Christensen. We conclude the paper with some possibly interesting questions.

We would like to thank the anonymous referee for his/her extremely careful reading of the paper and quite relevant suggestions.

1.4.1. Notation and conventions. The set of natural numbers is denoted either by \mathbb{N} or by \mathbb{Z}_+ . We denote by $\mathcal{M}(\mathbb{T})$ the space of all complex measures on \mathbb{T} , endowed with the total variation norm. The Fourier transform of a measure $\sigma \in \mathcal{M}(\mathbb{T})$ is denoted either by $\hat{\sigma}$ or by $\mathcal{F}(\sigma)$. As a rule, all measurable spaces (Ω, \mathfrak{A}) are standard Borel, and all measure spaces $(\Omega, \mathfrak{A}, m)$ are sigma-finite. All Hilbert spaces \mathcal{H} are

(complex) separable and infinite-dimensional. The scalar product $\langle u, v \rangle_{\mathcal{H}}$ is linear with respect to u and conjugate-linear with respect to v .

2. ABSTRACT RESULTS

2.1. S-mixing. It is well known (and easy to check) that the definitions of ergodicity, weak and strong mixing can be reformulated as follows. Let (X, \mathfrak{B}, μ) be a probability space, and set

$$L_0^2(\mu) := \left\{ f \in L^2(\mu); \int_X f d\mu = 0 \right\}.$$

Then, a measure-preserving transformation $T : (X, \mathfrak{B}, \mu) \rightarrow (X, \mathfrak{B}, \mu)$ is ergodic with respect to μ if and only if

$$\frac{1}{N} \sum_{n=0}^{N-1} \langle f \circ T^n, g \rangle_{L^2(\mu)} \xrightarrow{N \rightarrow \infty} 0$$

for any $f, g \in L_0^2(\mu)$; the transformation T is weakly mixing iff

$$\frac{1}{N} \sum_{n=0}^{N-1} |\langle f \circ T^n, g \rangle_{L^2(\mu)}| \xrightarrow{N \rightarrow \infty} 0$$

for any $f, g \in L_0^2(\mu)$; and T is strongly mixing iff

$$\langle f \circ T^n, g \rangle_{L^2(\mu)} \xrightarrow{n \rightarrow \infty} 0.$$

Now, let us denote by $V_T : L^2(\mu) \rightarrow L^2(\mu)$ the Koopman operator associated with a measure-preserving transformation $T : (X, \mathfrak{B}, \mu) \rightarrow (X, \mathfrak{B}, \mu)$, i.e. the isometry defined by

$$V_T f = f \circ T.$$

For any $f, g \in L^2(\mu)$, there is a uniquely defined complex measure $\sigma_{f,g} = \sigma_{f,g}^T$ on \mathbb{T} such that

$$\widehat{\sigma}_{f,g}(n) = \begin{cases} \langle V_T^n f, g \rangle_{L^2(\mu)} & \text{if } n \geq 0 \\ \langle V_T^{*|n|} f, g \rangle_{L^2(\mu)} & \text{if } n < 0 \end{cases}$$

(When $f = g$, this follows from Bochner's theorem because in this case the right-hand side defines a positive-definite function on \mathbb{Z} ; and then one can use a "polarization" argument). We denote by $\Sigma(T, \mu)$ the collection of all measures $\sigma_{f,g}$, $f, g \in L_0^2(\mu)$, and forgetting the measure μ we refer to $\Sigma(T, \mu)$ as "the spectral measure of T ".

With these notations, we see that T is weakly mixing with respect to μ iff all measures $\sigma \in \Sigma(T, \mu)$ are continuous (by Wiener's lemma), and that T is strongly mixing iff all measures $\sigma \in \Sigma(T, \mu)$ are Rajchman (by Rajchman's lemma). Likewise, T is ergodic iff $\sigma(\{\mathbf{1}\}) = 0$ for every $\sigma \in \Sigma(T, \mu)$.

More generally, given any family of measures $\mathcal{B} \subset \mathcal{M}(\mathbb{T})$, one may say that T is \mathcal{B} -mixing with respect to μ if the spectral measure of T lies in \mathcal{B} , i.e. all measures $\sigma \in \Sigma(T, \mu)$ are in \mathcal{B} . We shall in fact consider a more specific case which seems to

be the most natural one for our concerns. Let us denote by $\mathcal{F}_+ : \mathcal{M}(\mathbb{T}) \rightarrow \ell^\infty(\mathbb{Z}_+)$ the positive part of the Fourier transformation, i.e. $\mathcal{F}_+(\sigma) = \widehat{\sigma}|_{\mathbb{Z}_+}$.

Definition 2.1. Given any family $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$, we say that a measure $\sigma \in \mathcal{M}(\mathbb{T})$ is **S-continuous** if $\mathcal{F}_+(\sigma) \in \mathbf{S}$. A measure-preserving transformation $T : (X, \mu) \rightarrow (X, \mu)$ is **S-mixing** with respect to μ if every measure $\sigma \in \Sigma(T, \mu)$ is **S-continuous**. Accordingly, an operator $T \in \mathfrak{L}(X)$ is said to be **S-mixing in the Gaussian sense** if there exists a Gaussian, T -invariant measure μ on X with full support such that $T : (X, \mu) \rightarrow (X, \mu)$ is **S-mixing**.

Thus, strong mixing is just **S-mixing** for the family $\mathbf{S} = c_0(\mathbb{Z}_+)$, weak mixing is **S-mixing** for the the family \mathbf{S} of all sequences $(a_n) \in \ell^\infty(\mathbb{Z}_+)$ such that $|a_n| \rightarrow 0$ in the Cesàro sense, and ergodicity corresponds to the family \mathbf{S} of all $a \in \ell^\infty(\mathbb{Z}_+)$ tending to 0 in the Cesàro sense. In what follows, these families will be denoted by $\mathbf{S}_{\text{strong}}$, \mathbf{S}_{weak} and \mathbf{S}_{erg} , respectively.

2.2. Small subsets of the circle. Given a family of measures $\mathcal{B} \subset \mathcal{M}(\mathbb{T})$, it is quite natural in harmonic analysis to try to say something about the **B-small** subsets of \mathbb{T} , i.e. the sets $D \subset \mathbb{T}$ that are negligible for all positive measures $\sigma \in \mathcal{B}$. By this we mean that for any such measure σ , one can find a Borel set \widetilde{D} (possibly depending on σ) such that $D \subset \widetilde{D}$ and $\sigma(\widetilde{D}) = 0$. When the family \mathcal{B} has the form $\mathcal{B} = \mathcal{F}_+^{-1}(\mathbf{S})$ for some $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$, we call these sets **S-small**.

To avoid trivialities concerning **B-small** sets, the family \mathcal{B} under consideration should contain nonzero positive measures, and in fact it is desirable that it should be *hereditary* with respect to absolute continuity; that is, any measure absolutely continuous with respect to some $\sigma \in \mathcal{B}$ is again in \mathcal{B} . The following simple lemma shows how to achieve this for families of the form $\mathcal{F}_+^{-1}(\mathbf{S})$. Let us say that a family $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$ is *translation-invariant* if it is invariant under both the forward and the backward shift on $\ell^\infty(\mathbb{Z}_+)$.

Lemma 2.2. *If \mathbf{S} is a translation-invariant linear subspace of $\ell^\infty(\mathbb{Z}_+)$ such that $\mathcal{F}_+^{-1}(\mathbf{S})$ is norm-closed in $\mathcal{M}(\mathbb{T})$, then $\mathcal{F}_+^{-1}(\mathbf{S})$ is hereditary with respect to absolute continuity.*

Proof. If $\sigma \in \mathcal{F}_+^{-1}(\mathbf{S})$ then $P\sigma$ is in $\mathcal{F}_+^{-1}(\mathbf{S})$ for any trigonometric polynomial P , by translation-invariance. So the result follows by approximation. \square

We shall also make use of the following well known fact concerning **B-perfect** sets. By definition, a set $\Lambda \subset \mathbb{T}$ is **B-perfect** if $V \cap \Lambda$ is not **B-small** for any open set $V \subset \mathbb{T}$ such that $V \cap \Lambda \neq \emptyset$.

Lemma 2.3. *Let \mathcal{B} be a norm-closed linear subspace of $\mathcal{M}(\mathbb{T})$, and assume that \mathcal{B} is hereditary with respect to absolute continuity. For a closed set $\Lambda \subset \mathbb{T}$, the following are equivalent:*

- (a) Λ is **B-perfect**;
- (b) Λ is the support of some probability measure $\sigma \in \mathcal{B}$.

Proof. That (b) implies (a) is clear (without any assumption on \mathcal{B}). Conversely, assume that Λ is \mathcal{B} -perfect. Let $(W_n)_{n \geq 1}$ be a countable basis of open sets for Λ , with $W_n \neq \emptyset$. Since \mathcal{B} is hereditary, one can find for each n a probability measure $\sigma_n \in \mathcal{B}$ such that $\text{supp}(\sigma_n) \subset W_n$ and $\sigma_n(W_n) > 0$. Then the probability measure $\sigma = \sum_1^\infty 2^{-n} \sigma_n$ is in \mathcal{B} and $\text{supp}(\sigma) = \Lambda$. \square

2.3. The results. To state our results we need two more definitions.

Definition 2.4. Let T be an operator acting on a complex separable Fréchet space X , and let $\mathcal{B} \subset \mathcal{M}(\mathbb{T})$. We say that the \mathbb{T} -eigenvectors of T are \mathcal{B} -perfectly spanning if, for any Borel \mathcal{B} -small set $D \subset \mathbb{T}$, the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in X . When \mathcal{B} has the form $\mathcal{F}_+^{-1}(\mathbf{S})$, the terminology \mathbf{S} -perfectly spanning is used.

Thus, perfect spanning is \mathcal{B} -perfect spanning for the family of continuous measures, and \mathcal{U}_0 -perfect spanning is \mathcal{B} -perfect spanning for the family of Rajchman measures.

Remark. At some places, we will encounter sets which are *analytic* but perhaps not Borel. Recall that a set D in some Polish space Z is analytic if one can find a Borel relation $B \subset Z \times E$ (where E is Polish) such that $z \in D \Leftrightarrow \exists u : B(z, u)$. If the spanning property of the above definition holds for every analytic \mathcal{B} -small set D , we say that the \mathbb{T} -eigenvectors of T are \mathcal{B} -perfectly spanning *for analytic sets*.

Definition 2.5. We shall say that a family $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$ is c_0 -like if it has the form

$$\mathbf{S} = \{a \in \ell^\infty(\mathbb{Z}_+); \lim_{n \rightarrow \infty} \Phi_n(a) = 0\}$$

where $(\Phi_n)_{n \geq 0}$ is a uniformly bounded sequence of w^* -continuous semi-norms on $\ell^\infty(\mathbb{Z}_+)$. (By “uniformly bounded”, we mean that $\Phi_n(a) \leq C \|a\|_\infty$ for all n and some finite constant C).

For example, the families $\mathbf{S}_{\text{strong}}$, \mathbf{S}_{weak} and \mathbf{S}_{erg} are c_0 -like: just put $\Phi_n(a) = |a_n|$ in the strong mixing case, $\Phi_n(a) = \frac{1}{n} \sum_{k=0}^{n-1} |a_k|$ in the weak mixing case, and $\Phi_n(a) = \left| \frac{1}{n} \sum_{k=0}^{n-1} a_k \right|$ in the ergodic case.

Our main result is the following theorem, from which (1) and (2) in Theorem 1.1 follow immediately. Recall that a family $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$ is an *algebra* if it is a linear subspace which is closed under taking products (it is *not* assumed that \mathbf{S} contains $\mathbf{1}$), and that \mathbf{S} is *self-adjoint* if $\bar{u} \in \mathbf{S}$ whenever $u \in \mathbf{S}$.

Theorem 2.6. *Let X be a separable complex Fréchet space, and let $T \in \mathfrak{L}(X)$. Let also $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$. Assume that \mathbf{S} is a translation-invariant, self-adjoint and c_0 -like algebra, and that any \mathbf{S} -continuous measure is continuous. If the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning, then T is \mathbf{S} -mixing in the Gaussian sense.*

Remark. This theorem cannot be applied to the ergodic case, for two reasons: the family \mathbf{S}_{erg} is not an algebra, and \mathbf{S}_{erg} -continuous measures need not be continuous (a measure σ is \mathbf{S}_{erg} -continuous iff $\sigma(\{\mathbf{1}\}) = 0$).

Theorem 2.6 will be proved in section 5. The following much simpler converse result (which corresponds to (3) in Theorem 1.1) will be proved in section 6.

Proposition 2.7. *Let X be a separable complex Banach space, and assume that X has cotype 2. Let also \mathbf{S} be an arbitrary subset of $\ell^\infty(\mathbb{Z}_+)$. If $T \in \mathfrak{L}(X)$ is \mathbf{S} -mixing in the Gaussian sense, then the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning for analytic sets.*

Remark. Two “trivial” cases are worth mentioning. If we take $\mathbf{S} = \mathbf{S}_{\text{erg}}$, then $\mathcal{F}_+^{-1}(\mathbf{S}) = \{\sigma \in \mathcal{M}(\mathbb{T}); \sigma(\{\mathbf{1}\}) = 0\}$ and hence a set $D \subset \mathbb{T}$ is \mathbf{S} -small if and only if $D \subset \{\mathbf{1}\}$. If we take $\mathbf{S} = \ell^\infty(\mathbb{Z}_+)$, then $\mathcal{F}_+^{-1}(\mathbf{S}) = \mathcal{M}(\mathbb{T})$ and hence the only \mathbf{S} -small set is $D = \emptyset$. So, assuming that X has cotype 2, we get the following (known) results: if T admits an invariant Gaussian measure with full support, then the \mathbb{T} -eigenvectors of T span a dense subspace of X ; and if T is ergodic in the Gaussian sense, then $\overline{\text{span}} \left(\bigcup_{\lambda \in \mathbb{T} \setminus \{\mathbf{1}\}} \ker(T - \lambda) \right) = X$. As already pointed out, much more is true in the ergodic case: it is proved in [8, Theorem 4.1] that if T is ergodic in the Gaussian sense, then in fact the \mathbb{T} -eigenvectors are perfectly spanning (provided that X has cotype 2). Since ergodicity in the Gaussian sense is equivalent to weak mixing, this follows also from Proposition 2.7 applied with $\mathbf{S} = \mathbf{S}_{\text{weak}}$.

Our last result (to be proved also in section 6) says that when the space X is well-behaved, the assumptions in Theorem 2.6 can be relaxed: it is no longer necessary to assume that the family \mathbf{S} is c_0 -like, nor that \mathbf{S} -continuous measures are continuous. However, the (rather low) price to pay is that one has to use the strengthened version of \mathbf{S} -perfect spanning.

Theorem 2.8. *Let X be a separable complex Banach space, and assume that X has type 2. Let also \mathbf{S} be a norm-closed, translation-invariant and self-adjoint subalgebra of $\ell^\infty(\mathbb{Z}_+)$, and let $T \in \mathfrak{L}(X)$. If the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning for analytic sets, then T is \mathbf{S} -mixing in the Gaussian sense.*

Since Hilbert space has both type 2 and cotype 2, we immediately deduce

Corollary 2.9. *Let \mathbf{S} be a norm-closed, translation-invariant and self-adjoint subalgebra of $\ell^\infty(\mathbb{Z}_+)$. For Hilbert space operators, \mathbf{S} -mixing in the Gaussian sense is equivalent to the \mathbf{S} -spanning property of \mathbb{T} -eigenvectors (for analytic sets).*

Remark 1. The reader should wonder why analytic sets appear in Theorem 2.8, whereas only Borel sets are needed in Theorem 2.6. The reason is that the family of \mathbf{S} -small sets has a quite special structural property (the so-called *covering property*) when the family \mathbf{S} is c_0 -like: any analytic \mathbf{S} -small set can be covered by a sequence of closed \mathbf{S} -small sets (see [43]). In particular, any analytic \mathbf{S} -small set is contained in a Borel \mathbf{S} -small set and hence the two notions of \mathbf{S} -perfect spanning are equivalent.

In the weak mixing case, i.e. for the family of “small sets for continuous measures”, the covering property follows from the classical Perfect Set Theorem for analytic sets. In the strong mixing case, i.e. for the sets of extended uniqueness, this is a remarkable result due to G. Debs and J. Saint Raymond ([23]).

Remark 2. The proof of Theorem 2.8 turns out to be considerably simpler than that of Theorem 2.6. Roughly speaking, the reason is the following. Without any assumption on X , i.e. in the case of Theorem 2.6, we will have to be extremely careful to ensure that some “integral” operators of a certain kind are *gamma-radonifying* (see sub-section 3.1). On the other hand, such integral operators are automatically gamma-radonifying when X is a Banach space with type 2. So the main technical difficulty just disappears, and the proof becomes rather easy.

3. BACKGROUND

Throughout this section, X is a separable complex Fréchet space.

3.1. Gaussian measures and gamma-radonifying operators. This sub-section is devoted to a brief review of the basic facts that we shall need concerning Gaussian measures. For a reasonably self-contained exposition specifically tailored to linear dynamics, we refer to [9]; and for in-depth studies of Gaussian measures, to the books [13] and [17].

By a *Gaussian measure* on X , we mean a Borel probability measure μ on X which is the distribution of a random variable of the form

$$\sum_{n=0}^{\infty} g_n x_n$$

where (g_n) is a standard complex Gaussian sequence defined on some probability space $(\Omega, \mathfrak{A}, \mathbb{P})$ and (x_n) is a sequence in X such that the series $\sum g_n x_n$ is almost surely convergent.

We recall that (g_n) is a standard complex Gaussian sequence if the g_n are independent complex-valued random variables with distribution $\gamma_\sigma \otimes \gamma_\sigma$, where $\gamma_\sigma = \frac{1}{\sqrt{2\pi\sigma}} e^{-t^2/2\sigma^2} dt$ is the usual Gaussian distribution with mean 0 and variance $\sigma^2 = 1/2$. This implies in particular that $\mathbb{E}g_n = 0$ and $\mathbb{E}|g_n|^2 = 1$.

“Our” definition of a Gaussian measure is not the usual one. However, it is indeed equivalent to the classical definition, i.e. every continuous linear functional x^* has a complex symmetric Gaussian distribution when considered as a random variable on the probability space $(X, \mathfrak{B}(X), \mu)$ (where $\mathfrak{B}(X)$ is the Borel sigma-algebra of X).

It is well known that for a Fréchet space valued Gaussian series $\sum g_n x_n$, almost sure convergence is equivalent to convergence in the L^p sense, for any $p \in [1, \infty[$. It follows at once that if $\mu \sim \sum_{n \geq 0} g_n x_n$ is a Gaussian measure on X , then $\int_X \|x\|^2 d\mu(x) < \infty$ for every continuous semi-norm $\|\cdot\|$ on X . In particular, any continuous linear functional $x^* \in X^*$ is in $L^2(\mu)$ when considered as a random variable on (X, μ) . We also note that we consider only *centred* Gaussian measures μ , i.e. $\int_X x d\mu(x) = 0$.

Gaussian measures correspond in a canonical way to *gamma-radonifying operators*. Let \mathcal{H} be a Hilbert space. An operator $K : \mathcal{H} \rightarrow X$ is said to be gamma-radonifying if for some (equivalently, for any) orthonormal basis $(e_n)_{n \geq 0}$ of \mathcal{H} , the

Gaussian series $\sum g_n K(e_n)$ is almost surely convergent. A look at the Fourier transform reveals that the distribution of the random variable $\sum_{n \geq 0} g_n K(e_n)$ does not depend on the orthonormal basis (e_n) (see below); so the operator K gives rise to a Gaussian measure which may be denoted by μ_K :

$$\mu_K \sim \sum g_n K(e_n).$$

Conversely, it is not hard to show that any Gaussian measure $\mu \sim \sum g_n x_n$ is induced by some gamma-radonifying operator.

The following examples are worth keeping in mind: when X is a Hilbert space, an operator $K : \mathcal{H} \rightarrow X$ is gamma-radonifying if and only if it is a *Hilbert-Schmidt* operator; and for an arbitrary X , the operator K is gamma-radonifying as soon as $\sum_0^\infty \|K(e_n)\| < \infty$, for some orthonormal basis of \mathcal{H} and every continuous seminorm $\|\cdot\|$ on X . This follows at once from the equivalence between almost sure convergence and L^2 or L^1 convergence for Gaussian series.

We note that the support of a Gaussian measure $\mu \sim \sum_{n \geq 0} g_n x_n$ is the closed linear span of the x_n . Therefore, a Gaussian measure $\mu = \mu_K$ has full support if and only if the gamma-radonifying operator $K : \mathcal{H} \rightarrow X$ has dense range.

If $K : \mathcal{H} \rightarrow X$ is a continuous linear operator from some Hilbert space \mathcal{H} into X , we consider its adjoint K^* as an operator from X^* into \mathcal{H} . Hence, $K^* : X^* \rightarrow \mathcal{H}$ is a *conjugate-linear* operator. If K is gamma-radonifying and $\mu = \mu_K$ is the associated Gaussian measure, then we have the following fundamental identity:

$$\langle x^*, y^* \rangle_{L^2(\mu)} = \langle K^* y^*, K^* x^* \rangle_{\mathcal{H}}$$

for any $x^*, y^* \in X^*$. The proof is a simple computation using the orthogonality of the Gaussian variables g_n .

Let us also recall the definition of the *Fourier transform* of a Borel probability measure μ on X : this is the complex-valued function defined on the dual space X^* by

$$\widehat{\mu}(x^*) = \int_X e^{-i \operatorname{Re} \langle x^*, x \rangle} d\mu(x).$$

One very important property of the Fourier transform is that it uniquely determines the measure: if μ_1 and μ_2 have the same Fourier transform, then $\mu_1 = \mu_2$.

If $\mu \sim \sum_{n \geq 0} g_n x_n$ is a Gaussian measure, a simple computation shows that $\widehat{\mu}(x^*) = \exp\left(-\frac{1}{4} \sum_0^\infty |\langle x^*, x_n \rangle|^2\right)$ for any $x^* \in X^*$. It follows that if $K : \mathcal{H} \rightarrow X$ is a gamma-radonifying operator then

$$\widehat{\mu}_K(x^*) = e^{-\frac{1}{4} \|K^* x^*\|^2}.$$

In particular, two gamma-radonifying operators K_1 and K_2 define the same Gaussian measure if and only if

$$\|K_1^* x^*\| = \|K_2^* x^*\|$$

for every $x^* \in X^*$.

Finally, let us say a few words about type 2 and cotype 2 Banach spaces. A Banach space X has *type 2* if

$$\mathbb{E} \left\| \sum_n g_n x_n \right\| \leq C \left(\sum_n \|x_n\|^2 \right)^{1/2}$$

for some finite constant C and any finite sequence (x_n) in X ; and X has *cotype 2* if the reverse inequality holds. Thus, type 2 makes the convergence of Gaussian series easier, whereas on a cotype 2 space the convergence of such a series has strong implications. This is apparent in the following proposition, which contains all the results that we shall need concerning these notions. (See (3.1) below for the definition of the operators K_E).

Proposition 3.1. *Let X be a separable Banach space.*

- (1) *If X has type 2 then any operator $K_E : L^2(\Omega, m) \rightarrow X$ is gamma-radonifying.*
- (2) *If X has cotype 2 then any gamma-radonifying operator $K : L^2(\Omega, m) \rightarrow X$ has the form K_E , for some vector field $E : \Omega \rightarrow X$.*

These results are nontrivial (see e.g. [9]), but quite easy to use.

3.2. The general procedure. Most of what is needed for the the proof of Theorem 2.6 is summarized in Proposition 3.3 below, whose statement requires to introduce some terminology. Recall that all measure spaces under consideration are *sigma-finite*, and that all L^2 spaces are separable.

Let $(\Omega, \mathfrak{A}, m)$ be a measure space. By a (X -valued) *vector field* on $(\Omega, \mathfrak{A}, m)$ we mean any measurable map $E : \Omega \rightarrow X$ which is in $L^2(\Omega, m, X)$. Such a vector field gives rise to an operator $K_E : L^2(\Omega, m) \rightarrow X$ defined as follows:

$$(3.1) \quad K_E f = \int_{\Omega} f(\omega) E(\omega) dm(\omega).$$

It is easy to check that the operator K_E has dense range if and only if the vector field E is *m-spanning* in the following sense: for any measurable set $\Delta \subset \Omega$ with $m(\Delta) = 0$, the linear span of $\{E(\omega); \omega \in \Omega \setminus \Delta\}$ is dense in X . This happens in particular if Ω is a topological space, m is a Borel measure with full support, and the vector field E is continuous with $\overline{\text{span}} E(\Omega) = X$.

We also note that the operator K_E is always compact, and that it is a Hilbert-Schmidt operator if X is a Hilbert space (because E is in $L^2(\Omega, m, X)$). Moreover, the adjoint operator $K_E^* : X^* \rightarrow L^2(\Omega, m)$ is given by the formula

$$K_E^* x^* = \overline{\langle x^*, E(\cdot) \rangle}.$$

The next definition will be crucial for our purpose.

Definition 3.2. Let $T \in \mathfrak{L}(X)$. By a \mathbb{T} -*eigenfield* for T on $(\Omega, \mathfrak{A}, m)$ we mean a pair of maps (E, ϕ) where

- $E : \Omega \rightarrow X$ is a vector field;
- $\phi : \Omega \rightarrow \mathbb{T}$ is measurable;

- $TE(\omega) = \phi(\omega)E(\omega)$ for every $\omega \in \Omega$.

For example, if $E : \Lambda \rightarrow X$ is a continuous \mathbb{T} -eigenvector field for T defined on some compact set $\Lambda \subset \mathbb{T}$ and if we put $\phi(\lambda) = \lambda$, then (E, ϕ) is a \mathbb{T} -eigenfield for T on $(\Lambda, \mathfrak{B}(\Lambda), m)$ for any Borel probability measure m on Λ .

The key fact about \mathbb{T} -eigenfields is the following: if (E, ϕ) is a \mathbb{T} -eigenfield for T on (Ω, m) and if $M_\phi = L^2(\Omega, m) \rightarrow L^2(\Omega, m)$ is the (unitary) multiplication operator associated with ϕ , then the *intertwining equation*

$$TK_E = K_E M_\phi$$

holds. The proof is immediate.

We can now state

Proposition 3.3. *Let \mathbf{S} be a norm-closed, self-adjoint subalgebra of $\ell^\infty(\mathbb{Z}_+)$, and let $T \in \mathcal{L}(X)$. Assume that one can find a \mathbb{T} -eigenfield (E, ϕ) for T defined on some measure space (Ω, m) , such that*

- the operator $K_E : L^2(\Omega, m) \rightarrow X$ is gamma-radonifying;*
- the vector field E is m -spanning;*
- for any $f \in L^1(\Omega, m)$, the image measure $\sigma_f = (f m) \circ \phi^{-1}$ is \mathbf{S} -continuous.*

Then T is \mathbf{S} -mixing in the Gaussian sense. More precisely, T is \mathbf{S} -mixing with respect to μ_{K_E} .

As mentioned above, this proposition is essentially all what is needed to understand the proof of Theorem 2.6. It will follow at once from the next two lemmas, that will also be used in the proof of Proposition 2.7.

Lemma 3.4. *Let $T \in \mathcal{L}(X)$, and let $K : \mathcal{H} \rightarrow X$ be gamma-radonifying. Let also \mathbf{S} be a norm-closed, self-adjoint subalgebra of $\ell^\infty(\mathbb{Z}_+)$.*

- The measure $\mu = \mu_K$ is T -invariant if and only if one can find an operator $M : \mathcal{H} \rightarrow \mathcal{H}$ such that $\mathcal{H}_K := \mathcal{H} \ominus \ker(K)$ is M^* -invariant, M^* is an isometry on \mathcal{H}_K and $TK = KM$.*
- The operator T is \mathbf{S} -mixing with respect to μ if and only if $M^*_{|\mathcal{H}_K}$ has \mathbf{S} -continuous spectral measures, which means that the sequence $(\langle M^{*n}u, v \rangle_{\mathcal{H}})_{n \geq 0}$ is in \mathbf{S} for any $u, v \in \mathcal{H}_K$.*

Proof. Since T is a linear operator, the image measure $\mu_K \circ T^{-1}$ is equal to μ_{TK} . Taking the Fourier transforms, it follows that μ_K is T -invariant if and only if

$$\|K^*(x^*)\| = \|K^*(T^*x^*)\|$$

for every $x^* \in X^*$. This means exactly that one can find an isometry $V : \text{ran}(K^*) \rightarrow \text{ran}(K^*)$ such that $K^*T^* = VK^*$. Since $\overline{\text{ran}(K^*)} = \mathcal{H} \ominus \ker(K)$, this proves (1).

As for (2), the basic idea is very simple. Recall that T is \mathbf{S} -mixing with respect to μ if and only if

$$(3.2) \quad \mathcal{F}_+(\sigma_{f,g}) \in \mathbf{S}$$

for any $f, g \in L^2_0(\mu)$. When $f = x^*$ and $g = y^*$ are continuous linear functionals on X (recall that μ is centred, so that $X^* \subset L^2_0(\mu)$) we have $\widehat{\sigma}_{f,g}(n) = \langle f \circ T^n, g \rangle_{L^2(\mu)} =$

$\langle K^*T^{*n}(x^*), K^*y^* \rangle_{\mathcal{H}} = \langle M^{*n}(K^*x^*), K^*y^* \rangle_{\mathcal{H}}$ for every $n \geq 0$. Since M^* is power-bounded on \mathcal{H}_K and $\mathcal{H}_K = \text{ran}(K^*)$, it follows that $M^*|_{\mathcal{H}_K}$ has \mathbf{S} -continuous spectral measures if and only if (3.2) holds for all continuous linear functionals f, g . So the point is to pass from linear functionals to arbitrary $f, g \in L_0^2(\mu)$.

In the ‘‘classical’’ cases (weak and strong mixing), this can be done in at least two ways: either by reproducing a rather elementary argument of R. Rudnicki ([52, pp 108–109], see also [32] or [9, Theorem 5.24]), or by the more abstract approach of [7], which relies on the theory of *Fock spaces*. In the more general case we are considering, but assuming that \mathbf{S} is an *ideal* of $\ell^\infty(\mathbb{Z}_+)$, one possible proof would consist in merely copying out pp 5105–5108 of [7]. The fact that \mathbf{S} is norm-closed would be needed for the approximation argument on p. 5105, and the ideal property would be used on p. 5108 in the following way: if $(a_n) \in \mathbf{S}$ and if a sequence (b_n) satisfies $|b_n| \leq C|a_n|$ for all $n \in \mathbb{Z}_+$ (and some finite constant C), then $(b_n) \in \mathbf{S}$. Note that in the ‘‘classical’’ cases (weak mixing and strong mixing), the family \mathbf{S} is indeed an ideal of $\ell^\infty(\mathbb{Z}_+)$.

We outline a more direct approach, which is in fact exactly the same as in [7] but without any algebraic apparatus. In what follows, we denote by $\text{Re}(X^*)$ the set of all continuous, real-valued, real-linear functionals on X . For any $u \in \text{Re}(X^*)$, we denote by u^* the unique complex-linear functional with real part u , which is given by the formula $\langle u^*, x \rangle = u(x) - iu(ix)$.

First, we note that if (3.2) holds for all continuous linear functionals, then it holds for all $f, g \in \text{Re}(X^*)$. Indeed, using the invariance of μ under the transformation $x \mapsto ix$, it is easily checked that $\langle u, v \rangle_{L^2(\mu)} = \frac{1}{2} \text{Re}(\langle u^*, v^* \rangle_{L^2(\mu)})$ for any $u, v \in \text{Re}(X^*)$. Applying this with $u := f \circ T^n$, $n \geq 0$ and $v := g$ and since \mathbf{S} is closed under taking real parts, it follows that if $f, g \in \text{Re}(X^*)$ and $\mathcal{F}_+(\sigma_{f^*, g^*}) \in \mathbf{S}$ then $\mathcal{F}_+(\sigma_{f, g}) \in \mathbf{S}$.

Now, let us denote by H_k , $k \geq 0$ the classical real *Hermite polynomials*, i.e. the orthogonal polynomials associated with the standard Gaussian distribution $\gamma = \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$ on \mathbb{R} . For every $k \geq 0$, set

$$\mathcal{H}_k := \overline{\text{span}}^{L^2(\mu)} \{H_k(f); f \in \mathcal{S}\},$$

where \mathcal{S} is the set of all $f \in \text{Re}(X^*)$ such that $\|f\|_{L^2(\mu)} = 1$. It is well known that $L^2(\mu)$ is the orthogonal direct sum of the subspaces \mathcal{H}_k , $k \geq 0$ (this is the so-called *Wiener chaos decomposition*) and hence that $L_0^2(\mu) = \bigoplus_{k \geq 1} \mathcal{H}_k$. Moreover, it is also well known that

$$(3.3) \quad \langle H_k(u), H_k(v) \rangle_{L^2(\mu)} = \langle u, v \rangle_{L^2(\mu)}^k$$

for any $u, v \in \mathcal{S}$ and every $k \geq 0$ (see e.g. [21, Chapter 9], where the proofs are given in a Hilbert space setting but work equally well on a Fréchet space). Taking $u := f \circ T^n$, $n \geq 0$ and $v := g$, it follows that $\mathcal{F}_+(\sigma_{H_k(f), H_k(g)}) = [\mathcal{F}_+(\sigma_{f, g})]^k$ for any $f, g \in \mathcal{S}$ and all $k \geq 0$. Since \mathbf{S} is a closed subalgebra of $\ell^\infty(\mathbb{Z}_+)$ and since the map $(f, g) \mapsto \mathcal{F}_+(\sigma_{f, g})$ is continuous from $L^2(\mu) \times L^2(\mu)$ into $\ell^\infty(\mathbb{Z}_+)$ we conclude that (3.2) does hold true for any $f, g \in L_0^2(\mu)$ as soon as it holds for linear functionals. \square

Remark 1. It is apparent from the above proof that the implication “ T is \mathbf{S} -mixing implies $M_{|\mathcal{H}_K}^*$ has \mathbf{S} -continuous spectral measures” requires no assumption on the family \mathbf{S} . This will be used in the proof of Proposition 2.7.

Remark 2. One can use (3.3) to recover the well-known fact that ergodicity in the Gaussian sense is equivalent to weak mixing. Indeed, if T is ergodic with respect to the Gaussian measure μ then, applying (3.3) with $k = 2$ and using ergodicity, we see that

$$\frac{1}{N} \sum_{n=0}^{N-1} \langle u \circ T^n, v \rangle_{L^2(\mu)}^2 = \frac{1}{N} \sum_{n=0}^{N-1} \langle \mathcal{H}_2(u) \circ T^n, \mathcal{H}_2(v) \rangle_{L^2(\mu)} \xrightarrow{N \rightarrow \infty} 0$$

for any $u, v \in \mathcal{S}$. As is well known, the same property then holds with $\langle u \circ T^n, v \rangle_{L^2(\mu)}^2$ replaced with $|\langle u \circ T^n, v \rangle_{L^2(\mu)}|$, and hence we get the weak mixing property.

Lemma 3.5. *Let $M = M_\phi$ be a unitary multiplication operator on $\mathcal{H} = L^2(\Omega, m)$ associated with a measurable function $\phi : \Omega \rightarrow \mathbb{T}$. Let also $\mathcal{H}_1 \subset \mathcal{H}$ be a closed M^* -invariant subspace, and let us denote by $\mathcal{H}_1 \cdot \mathcal{H}$ the set of all $f \in L^1(m)$ that can be written as $f = h_1 h$, where $h_1 \in \mathcal{H}_1$ and $h \in \mathcal{H}$. Finally, let \mathbf{S} be an arbitrary subset of $\ell^\infty(\mathbb{Z}_+)$. Consider the following assertions:*

- (i) $M_{|\mathcal{H}_1}^*$ has \mathbf{S} -continuous spectral measures;
- (ii) for any $f \in \mathcal{H}_1 \cdot \mathcal{H}$, the image measure $\sigma_f = (fm) \circ \phi^{-1}$ is \mathbf{S} -continuous;
- (iii) $\mathbf{1}_{\{\phi \in D\}} h \perp \mathcal{H}_1$, for any \mathbf{S} -small analytic set $D \subset \mathbb{T}$ and every $h \in \mathcal{H}$.

Then (i) and (ii) are equivalent and imply (iii).

Proof. We note that (iii) makes sense because $\phi^{-1}(D)$ is m -measurable for any analytic $D \subset \mathbb{T}$ (see [39]).

A straightforward computation shows that for any $u, v \in L^2(\Omega, m)$, the Fourier coefficients of $\sigma_{u\bar{v}}$ are given by

$$\hat{\sigma}_{u\bar{v}}(n) = \langle M^{*n}u, v \rangle_{\mathcal{H}}.$$

Moreover, since \mathcal{H}_1 is M^* -invariant we have

$$\langle M^{*n}h_1, h \rangle_{\mathcal{H}} = \langle M^{*n}h_1, \pi_1 h \rangle_{\mathcal{H}} \quad (n \geq 0)$$

for any $h_1 \in \mathcal{H}_1$ and every $h \in \mathcal{H}$, where π_1 is the orthogonal projection onto \mathcal{H}_1 . It follows that $M_{|\mathcal{H}_1}^*$ has \mathbf{S} -continuous spectral measures if and only if $\mathcal{F}_+(\sigma_f) \in \mathbf{S}$ for any $f = h_1 \bar{h} \in \mathcal{H}_1 \cdot \mathcal{H}$. In other words, (i) and (ii) are equivalent.

For any analytic set $D \subset \mathbb{T}$ and $h, h_1 \in \mathcal{H}$, we have

$$\langle h_1, \mathbf{1}_{\{\phi \in D\}} h \rangle_{\mathcal{H}} = \sigma_{h_1 \bar{h}}(D).$$

Hence, (iii) says exactly that $\sigma_f(D) = 0$ for any $f \in \mathcal{H}_1 \cdot \mathcal{H}$ and every \mathbf{S} -small analytic set $D \subset \mathbb{T}$. From this, it is clear that (ii) implies (iii). \square

Remark. Properties (i), (ii), (iii) are in fact equivalent in the strong mixing case, i.e. $\mathbf{S} = c_0(\mathbb{Z}_+)$. Indeed, by a famous theorem of R. Lyons ([42], see also [40]), a positive measure σ is Rajchman if and only if $\sigma(D) = 0$ for every Borel set of extended

uniqueness $D \subset \mathbb{T}$. From this, it follows that if (iii) holds then σ_f is Rajchman for every nonnegative $f \in \mathcal{H} \cdot \mathcal{H}_1$, and it is easy to check that this implies (ii).

Proof of Proposition 3.3. Let $M_\phi : L^2(\Omega, m) \rightarrow L^2(\Omega, m)$ be the unitary multiplication operator associated with ϕ . Since $TK_E = K_E M_\phi$, the Gaussian measure $\mu = \mu_{K_E}$ is T -invariant by Lemma 3.4. Moreover, μ has full support since K_E has dense range (by (b)), and T is \mathbf{S} -mixing with respect to μ by Lemmas 3.4 and 3.5. \square

Remark 1. An examination of the proof reveals that assumption (c) in Proposition 3.3 is a little bit stronger than what is actually needed: since $K_E^* : X^* \rightarrow L^2(\Omega, m)$ is given by $K_E^* x^* = \overline{\langle x^*, E(\cdot) \rangle}$, it is enough to assume that the measure $(fm) \circ \phi^{-1}$ is \mathbf{S} -continuous for every $f \in L^1(\Omega, m)$ of the form $f = \langle x^*, E(\cdot) \rangle \overline{\langle y^*, E(\cdot) \rangle}$, where $x^*, y^* \in X^*$. However, the proposition is easier to remember as stated.

Remark 2. Somewhat ironically, it will follow from Theorem 2.6 that the seemingly crucial gamma-radonifying assumption (a) in Proposition 3.3 is in fact not necessary to ensure \mathbf{S} -mixing in the Gaussian sense. Indeed, it is easily checked that if one can find a \mathbb{T} -eigenfield for T satisfying (b) and (c), then the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning. In fact, the proof of Theorem 2.6 essentially consists in showing that if there exists a \mathbb{T} -eigenfield satisfying (b) and (c), then it is possible to construct one satisfying (b), (c) and (a).

3.3. An “exhaustion” lemma. Besides Proposition 3.3, the following lemma will also be needed in the proof of Theorems 2.6 and 2.8.

Lemma 3.6. *Let $\mathcal{B} \subset \mathcal{M}(\mathbb{T})$, and let $T \in \mathcal{L}(X)$. Assume that the \mathbb{T} -eigenvectors of T are \mathcal{B} -perfectly spanning for analytic sets. Finally, put*

$$\mathbf{V} := \{(x, \lambda) \in X \times \mathbb{T}; x \neq 0 \text{ and } T(x) = \lambda x\}.$$

Then, one can find a closed subset \mathbf{Z} of \mathbf{V} with the following properties:

- *for any (relatively) open set $O \subset \mathbf{V}$ such that $O \cap \mathbf{Z} \neq \emptyset$, the set $\pi_2(O \cap \mathbf{Z}) := \{\lambda \in \mathbb{T}; \exists x \in O \cap \mathbf{Z} : (x, \lambda) \in \mathbf{Z}\}$ is not \mathcal{B} -small;*
- *the linear span of $\pi_1(\mathbf{Z}) := \{x \in X; \exists \lambda \in \mathbb{T} : (x, \lambda) \in \mathbf{Z}\}$ is dense in X .*

Proof. Let us denote by \mathbf{O} the union of all relatively open sets $O \subset \mathbf{V}$ such that $\pi_2(O)$ is \mathcal{B} -small, and set $\mathbf{Z} := \mathbf{V} \setminus \mathbf{O}$. Then \mathbf{Z} is closed in \mathbf{V} and satisfies the first required property (by its very definition). Moreover, by the Lindelöf property and since any countable union of \mathcal{B} -small sets is \mathcal{B} -small, the set $D := \pi_2(\mathbf{O})$ is \mathcal{B} -small; and D is an analytic set because \mathbf{O} is Borel in $X \times \mathbb{T}$. By assumption on T , the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in X . Now, any \mathbb{T} -eigenvector x for T is in $\pi_1(\mathbf{V})$, and if the associated eigenvalue λ is not in D then $(x, \lambda) \in \mathbf{Z}$ by the definition of D . It follows that $\ker(T - \lambda) \setminus \{0\}$ is contained in $\pi_1(\mathbf{Z})$ for any $\lambda \in \mathbb{T} \setminus D$, and hence that $\overline{\text{span}}(\pi_1(\mathbf{Z})) = X$. \square

Despite its simplicity, Lemma 3.6 will be quite useful for us. We illustrate it by proving the following result.

Proposition 3.7. *Let $\mathcal{B} \subset \mathcal{M}(\mathbb{T})$ be hereditary with respect to absolute continuity, and let $T \in \mathfrak{L}(X)$. The following are equivalent:*

- (i) *the \mathbb{T} -eigenvectors of T are \mathcal{B} -perfectly spanning for analytic sets;*
- (ii) *one can find a countable family of continuous \mathbb{T} -eigenvector fields $(E_i)_{i \in I}$ for T , where $E_i : \Lambda_i \rightarrow X$ is defined on some \mathcal{B} -perfect set $\Lambda_i \subset \mathbb{T}$, such that $\text{span}(\bigcup_{i \in I} E_i(\Lambda_i))$ is dense in X .*

Proof. Since the E_i are assumed to be continuous in (ii), it is plain that (ii) implies (i). Conversely, assume that (i) holds true, and let \mathbf{Z} be the set given by Lemma 3.6.

Choose a sequence $(x_i, \lambda_i)_{i \in \mathbb{N}} \subset \mathbf{Z}$ such that the set $\{(x_i, \lambda_i); i \in I\}$ is dense in \mathbf{Z} for every cofinite set $I \subset \mathbb{N}$, and let $(\varepsilon_i)_{i \in \mathbb{N}}$ be any sequence of positive numbers tending to 0. Let also d be a compatible metric d on X and put

$$B_i := \{x \in X; d(x, x_i) < \varepsilon_i\}.$$

By the definition of \mathbf{Z} and since \mathcal{B} is hereditary with respect to absolute continuity, one can find for each $i \in \mathbb{N}$, a probability measure $\sigma'_i \in \mathcal{B}$ such that

$$\text{supp}(\sigma'_i) \subset \{\lambda \in \mathbb{T}; \exists x \in B_i : T(x) = \lambda x\}.$$

Put $\Lambda'_i := \text{supp}(\sigma'_i)$. The set

$$A_i = \{(x, \lambda) \in B_i \times \Lambda'_i; T(x) = \lambda x\}$$

is closed in the Polish space $B_i \times \Lambda'_i$ and projects onto Λ'_i . By the Jankov–von Neumann uniformization theorem (see [39, 18.A]), one can find a universally measurable map $E'_i : \Lambda'_i \rightarrow B_i$ such that $(E'_i(\lambda), \lambda) \in A_i$ for every $\lambda \in \Lambda'_i$. In other words we have a universally measurable \mathbb{T} -eigenvector field $E'_i : \Lambda'_i \rightarrow X$ such that $d(E'_i(\lambda), x_i) < \varepsilon_i$ for every $\lambda \in \Lambda'_i$. By Lusin's theorem on measurable functions (see e.g. [39, 17.D]), one can find a closed set C_i of positive σ'_i -measure such that $(E'_i)|_{C_i}$ is continuous. Then the measure $\sigma_i = (\sigma'_i)|_{C_i}$ is in \mathcal{B} since it is absolutely continuous with respect to σ_i , so the set $\Lambda_i = \text{supp}(\sigma_i)$ is \mathcal{B} -perfect. The map $E_i := (E'_i)|_{\Lambda_i} : \Lambda_i \rightarrow X$ is continuous, and we have $d(E_i(\lambda), x_i) < \varepsilon_i$ for every $\lambda \in \Lambda_i$. Since $\text{span}\{x_i; i \in I\}$ is dense in X for every cofinite set $I \subset \mathbb{N}$ by Lemma 3.6, it follows that $\text{span}(\bigcup_{i \in \mathbb{N}} E_i(\Lambda_i))$ is dense in X , which concludes the proof. \square

Remark. When \mathcal{B} is the family of continuous measures, the equivalence of (i) and (ii) is proved in [30, Proposition 4.2].

4. THE WEAK MIXING CASE

In this section, we concentrate on part (1) of Theorem 1.1. So we assume that the \mathbb{T} -eigenvectors of our operator $T \in \mathfrak{L}(X)$ are perfectly spanning, and we want to show that T is weakly mixing in the Gaussian sense. For simplicity, we assume that X is a *Banach* space in order to avoid working with sequences of continuous semi-norms.

4.1. Why things may not be obvious. Before embarking on the proof, let us briefly explain why it could go wrong. In view of Proposition 3.3, we need to construct a \mathbb{T} -eigenfield (E, ϕ) for T on some measure space (Ω, m) such that

- (a) the operator $K_E : L^2(\Omega, m) \rightarrow X$ is gamma-radonifying;
- (b) the vector field E is m -spanning;
- (c) for any $f \in L^1(\Omega, m)$, the image measure $\sigma_f = (f m) \circ \phi^{-1}$ is continuous.

To achieve (a) and (b), the most brutal way is to choose some total sequence $(x_n)_{n \in \mathbb{N}} \subset B_X$ consisting of \mathbb{T} -eigenvectors for T , say $T(x_n) = \lambda_n x_n$, to take $\Omega = \mathbb{N}$ with the counting measure m , and to define (E, ϕ) by $E(n) = 2^{-n} x_n$ and $\phi(n) = \lambda_n$. Then the vector field E is obviously m -spanning, and the operator K_E is gamma-radonifying since $K_E(e_n) = 2^{-n} x_n$ and hence $\sum_0^\infty \|K_E(e_n)\| < \infty$. (Here, of course, (e_n) is the canonical basis of $L^2(\Omega, m) = \ell^2(\mathbb{N})$). However, the image measure $\sigma = m \circ \phi^{-1}$ is purely atomic, so (c) is certainly not satisfied. (In fact, by [30, Theorem 5.1], T will never be ergodic with respect to any measure induced by a random variable of the form $\xi = \sum_0^\infty \chi_n x_n$, where (x_n) is a sequence of \mathbb{T} -eigenvectors for T and (χ_n) is a sequence of independent, rotation-invariant centred random variables). To get (c) we must take a more complicated measure space (Ω, m) and avoid atoms; but then it will be harder to show that an operator defined on $L^2(\Omega, m)$ is gamma-radonifying. Thus, conditions (a) and (c) go in somewhat opposite directions, and we have to find a kind of balance between them.

In [8] and [9], the measure space (Ω, m) was an open arc of \mathbb{T} (with the Lebesgue measure), and the difficulty was partially settled by requiring some regularity on the \mathbb{T} -eigenvector fields and combining this with the geometrical properties of the underlying Banach space. In the present paper, we will allow more freedom on the measure space, and this will enable us to get rid of any assumption on X .

The main part of the proof is divided into two steps. We shall first explain how to produce gamma-radonifying operators of the form K_E in a reasonably flexible way, and then we show how this can be used to construct suitable \mathbb{T} -eigenfields for T under the perfectly spanning assumption. Once this has been done, the conclusion follows easily.

4.2. How to construct gamma-radonifying operators. The first part of our program relies essentially on the following observation. Let G be a compact metrizable abelian group with normalized Haar measure m_G and dual group Γ , and let $E : G \rightarrow X$ be a vector field on (G, m_G) . Then the operator $K_E : L^2(G, m_G) \rightarrow X$ is gamma-radonifying as soon as

$$\sum_{\gamma \in \Gamma} \|\widehat{E}(\gamma)\| < \infty,$$

where \widehat{E} is the Fourier transform of E . This is indeed obvious since the characters of G form an orthonormal basis $(e_\gamma)_{\gamma \in \Gamma}$ of $L^2(G, m_G)$ and $K_E(e_\gamma) = \widehat{E}(\gamma)$ for every $\gamma \in \Gamma$.

The compact group we shall use is the usual *Cantor group* $G := \{0, 1\}^{\mathbb{N}}$, where addition is performed modulo 2. The dual group Γ is identified with the set all of

finite subsets of \mathbb{N} , which we denote by FIN . A set $I \in \text{FIN}$ corresponds to the character $\gamma_I \in \Gamma$ defined by the formula

$$\gamma_I(\omega) = \prod_{i \in I} \varepsilon_i(\omega),$$

where $\varepsilon_i(\omega_0, \omega_1, \dots) = (-1)^{\omega_i}$. An empty product is declared to be equal to 1, so that $\gamma_\emptyset = \mathbf{1}$.

It is common knowledge that any ‘‘sufficiently regular’’ function has an absolutely convergent Fourier series. In our setting, regularity will be quantified as follows. Let us denote by d the usual ultrametric distance on G ,

$$d(\omega, \omega') = 2^{-n(\omega, \omega')},$$

where $n(\omega, \omega')$ is the first i such that $\omega_i \neq \omega'_i$. We shall say that a map $E : G \rightarrow X$ is *super-Lipschitz* if it is α -Hölderian for some $\alpha > 1$, i.e.

$$\|E(\omega) - E(\omega')\| \leq C d(\omega, \omega')^\alpha$$

for any $\omega, \omega' \in G$ and some finite constant C . (Of course, there are no nonconstant super-Lipschitz maps on \mathbb{T} ; but life is different on the Cantor group).

The following lemma is the kind of result that we need.

Lemma 4.1. *If $E : G \rightarrow X$ is super-Lipschitz, then E has an absolutely convergent Fourier series.*

Proof. Assume that E is α -Hölderian, $\alpha > 1$, with constant C . The key point is the following

Fact. Let $n \in \mathbb{N}$. If $I \in \text{FIN}$ satisfies $I \ni n$ then $\|\widehat{E}(\gamma_I)\| \leq (C/2) \times 2^{-\alpha n}$.

Indeed, setting $\mathcal{F}_n := \{I \in \text{FIN}; I \neq \emptyset \text{ and } \max I = n\}$ (which has cardinality 2^n), this yields that

$$\sum_{I \in \mathcal{F}_n} \|\widehat{E}(\gamma_I)\| \leq 2^n \times (C/2) \times 2^{-\alpha n} = (C/2) \times 2^{-\beta n}$$

for all $n \geq 0$ (where $\beta = \alpha - 1 > 0$), and the result follows.

Proof of Fact. Write $I = J \cup \{n\}$, where $J \in \text{FIN}$ and $n \notin J$. Then

$$\langle \gamma_I, \omega \rangle = \varepsilon_n(\omega) \langle \gamma_J, \omega \rangle,$$

and hence

$$\begin{aligned} \widehat{E}(\gamma_I) &= \int_{\{\omega; \varepsilon_n(\omega)=1\}} \langle \gamma_J, \omega \rangle E(\omega) dm_G(\omega) - \int_{\{\omega; \varepsilon_n(\omega)=-1\}} \langle \gamma_J, \omega \rangle E(\omega) dm_G(\omega) \\ &= \int_{\{\omega; \varepsilon_n(\omega)=1\}} \langle \gamma_J, \omega \rangle [E(\omega) - E(\omega + \delta^n)] dm_G(\omega), \end{aligned}$$

where $\delta^n \in G$ is the sequence which is 0 everywhere except at the n -th coordinate. By assumption on E , we know that $\|E(\omega) - E(\omega + \delta^n)\| \leq C \times 2^{-\alpha n}$; and since the set $\{\varepsilon_n = 1\}$ has measure $1/2$, this concludes the proof. \square

\square

4.3. Construction of suitable \mathbb{T} -eigenfields. The second part of our program is achieved by the following lemma.

Lemma 4.2. *Put $\mathbf{V} := \{(x, \lambda) \in X \times \mathbb{T}; x \neq 0 \text{ and } T(x) = \lambda x\}$, and let $\mathbf{Z} \subset \mathbf{V}$. Assume that for any (relatively) open set $O \subset \mathbf{V}$ such that $O \cap \mathbf{Z} \neq \emptyset$, the set $\{\lambda \in \mathbb{T}; \exists x : (x, \lambda) \in O\}$ is uncountable. Then, for any (x_0, λ_0) in \mathbf{Z} and $\varepsilon > 0$, one can construct a \mathbb{T} -eigenfield (E, ϕ) for T on the Cantor group (G, m_G) such that*

- (1) $\|E(\omega) - x_0\| < \varepsilon$ for all $\omega \in G$;
- (2) $E : G \rightarrow X$ is super-Lipschitz;
- (3) $\phi : G \rightarrow \mathbb{T}$ is a homeomorphic embedding.

Proof. This is a standard Cantor-like construction. Let us denote by \mathcal{S} the set of all finite 0-1 sequences (including the empty sequence \emptyset). We write $|s|$ for the length of a sequence $s \in \mathcal{S}$, and if $i \in \{0, 1\}$ we denote by si the sequence “ s followed by i ”.

Put $V_\emptyset := \mathbb{T}$ and choose an open set $U_\emptyset \subset X$ with $x_0 \in U_\emptyset$ and $\text{diam}(U_\emptyset) < \varepsilon$. Let us also fix an arbitrary Hölder exponent $\alpha > 1$. We construct inductively two sequences of open sets $(U_s)_{s \in \mathcal{S}}$ in X and $(V_s)_{s \in \mathcal{S}}$ in \mathbb{T} , such that the following properties hold for all $s \in \mathcal{S}$ and $i \in \{0, 1\}$.

- (i) $\overline{U_{si}} \subset U_s$ and $\overline{V_{si}} \subset V_s$;
- (ii) $\overline{U_{s0}} \cap \overline{U_{s1}} = \emptyset$ and $\overline{V_{s0}} \cap \overline{V_{s1}} = \emptyset$;
- (iii) $\text{diam}(U_s) \leq 2^{-\alpha|s|}$ and $\text{diam}(V_{si}) \leq \frac{1}{2} \text{diam}(V_s)$;
- (iv) $(U_s \times V_s) \cap \mathbf{Z} \neq \emptyset$.

The inductive step is easy. Assume that U_s and V_s have been constructed for some s . Since $(U_s \times V_s) \cap \mathbf{Z} \neq \emptyset$, we know that the set $\{\lambda \in \mathbb{T}; \exists x : (x, \lambda) \in (U_s \times V_s) \cap \mathbf{Z}\}$ is uncountable, and hence contains at least 2 distinct points λ^0, λ^1 . Pick $x^i \in U_s$ such that $(x^i, \lambda^i) \in \mathbf{Z}$. Then $x^0 \neq x^1$ because $T(x^i) = \lambda^i x^i$ and $x^i \neq 0$. Thus, one may choose small enough open sets $U_{si} \ni x^i$ and $V_{si} \ni \lambda^i$ to ensure (i), ..., (iv) at steps $s0$ and $s1$.

For any $\omega \in G$ and $n \in \mathbb{N}$, let us denote by $\omega|_n$ the finite sequence $(\omega_0, \dots, \omega_n) \in \mathcal{S}$. By (i), (ii), (iii), the intersection $\bigcap_{n \geq 0} U_{\omega|_n}$ is a single point $\{x_\omega\}$, and the intersection $\bigcap_{n \geq 0} V_{\omega|_n}$ is a single point $\{\lambda_\omega\}$. Moreover, the map $\omega \mapsto \lambda_\omega := \phi(\omega)$ is a homeomorphic embedding, and the map $\omega \mapsto x_\omega := E(\omega)$ is α -Hölderian and hence super-Lipschitz (by (iii)). Finally, it follows easily from the continuity of T that $T(x_\omega) = \lambda_\omega x_\omega$ for every $\omega \in G$. In other words, (E, ϕ) is a \mathbb{T} -eigenfield for T . Since $\|E(\omega) - x_0\| \leq \text{diam}(U_\emptyset) < \varepsilon$ for all $\omega \in G$, this concludes the proof. \square

4.4. The proof. We now just have to put the pieces together. Let us recall what we are trying to do: we are given an operator $T \in \mathfrak{L}(X)$ whose \mathbb{T} -eigenvectors are perfectly spanning, and we want to show that T is weakly mixing in the Gaussian sense.

Let $\mathbf{V} := \{(x, \lambda) \in X \times \mathbb{T}; x \neq 0 \text{ and } T(x) = \lambda x\}$. Applying Lemma 3.6 to the family of continuous measures, we get a closed set $\mathbf{Z} \subset \mathbf{V}$ with the following properties:

- \mathbf{Z} satisfies the assumption of Lemma 4.2, i.e. for any $O \subset \mathbf{V}$ open such that $O \cap \mathbf{Z} \neq \emptyset$, the set $\{\lambda \in \mathbb{T}; \exists x : (x, \lambda) \in O\}$ is uncountable;
- $\text{span}(\pi_1(\mathbf{Z}))$ is dense in X .

Let us fix a sequence $(x_i, \lambda_i)_{i \in \mathbb{N}} \subset \mathbf{Z}$ such that the set $\{(x_i, \lambda_i); i \in I\}$ is dense in \mathbf{Z} for every cofinite set $I \subset \mathbb{N}$, and let us apply Lemma 4.2 to each point (x_i, λ_i) , with e.g. $\varepsilon_i = 2^{-i}$. Taking Lemma 4.1 into account and since $\varepsilon_i \rightarrow 0$, this give a sequence of \mathbb{T} -eigenfields (E_i, ϕ_i) defined on $(\Omega_i, m_i) := (G, m_G)$, such that

- (1) each operator $K_{E_i} : L^2(\Omega_i, m_i) \rightarrow X$ is gamma-radonifying;
- (2) each E_i is continuous and $\overline{\text{span}}(\bigcup_{i \in \mathbb{N}} E_i(\Omega_i)) = X$;
- (3) each ϕ_i is one-to-one.

Now, let (Ω, m) be the “disjoint union” of the measure spaces (Ω_i, m_i) (so that $L^2(\Omega, m)$ is the ℓ^2 -direct sum $\bigoplus_i L^2(\Omega_i, m_i)$). Choose a sequence of small positive numbers $(\alpha_i)_{i \in \mathbb{N}}$, and define a \mathbb{T} -eigenfield (E, ϕ) on (Ω, m) as follows: $E(\omega_i) = \alpha_i E_i(\omega_i)$ and $\phi(\omega_i) = \phi_i(\omega_i)$ for each i and every $\omega_i \in \Omega_i$. If the α_i are small enough, then E is indeed in $L^2(\Omega, m)$ and (using property (1)) the operator $K_E : L^2(\Omega, m) \rightarrow X$ is gamma-radonifying. By (2) and since m has full support, the vector field E is m -spanning and hence the operator K_E has dense range. Moreover, since $TK_{E_i} = K_{E_i}M_{\phi_i}$ for all i , the intertwining equation $TK_E = K_E M_\phi$ holds. Finally, since (Ω_i, m_i) is atomless, it follows from (3) that each measure $m_i \circ \phi_i^{-1}$ is continuous, and hence that $\sigma = m \circ \phi^{-1}$ is continuous as well. By Proposition 3.3, we conclude that T is weakly mixing in the Gaussian sense.

5. PROOF OF THE ABSTRACT RESULTS (1)

In this section, we prove Theorem 2.6. The proof runs along exactly the same lines as that of the weak mixing case: we first explain how to produce gamma-radonifying operators of the form K_E in a flexible way, then we show how this can be used to construct suitable \mathbb{T} -eigenfields for T under the \mathbf{S} -perfectly spanning assumption, and the conclusion follows easily. However, the first two steps are technically more involved than the corresponding ones from the weak mixing case, the difficulty being not less important when dealing with the strong mixing case only. Roughly speaking, the main reason is that it is much harder for a measure to be Rajchman, or more generally to be \mathbf{S} -continuous, than to be merely continuous.

To avoid artificial complications, we shall first assume that the underlying space X is a Banach space (which will dispense us from using sequences of semi-norms), and we will indicate only at the very end of the proof how it can be adapted in the Fréchet space case (this is really a matter of very minor changes). Thus, in most of this section, T is a linear operator acting on a complex separable Banach space X . We are also given a norm-closed, c_0 -like, translation-variant and self-adjoint algebra $\mathbf{S} \subset \ell^\infty(\mathbb{Z}_+)$ such that every \mathbf{S} -continuous measure is continuous. We are assuming that the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning, and our aim is to show that T is \mathbf{S} -mixing in the Gaussian sense.

5.1. How to construct gamma-radonifying operators. Just as in the weak mixing case, the guiding idea of the first part of our program is the observation

that we made in sub-section 4.2, namely that if $E : \Omega \rightarrow X$ is a vector field defined on a compact abelian group Ω (with Haar measure m and dual group Γ), then the operator $K_E : L^2(\Omega, m) \rightarrow X$ is gamma-radonifying as soon as

$$\sum_{\gamma \in \Gamma} \|\widehat{E}(\gamma)\| < \infty.$$

Unfortunately, we are not able to use this observation as stated. Instead of compact groups, we will be forced to use some slightly more complicated objects (Ω, m) , due to the structure of the inductive construction that will be performed in the *second part* of our program.

Let us denote by \mathfrak{Q} the set of all finite sequences of integers $\bar{q} = (q_1, \dots, q_l)$ with $q_s \geq 2$ for all s . For any $\bar{q} = (q_1, \dots, q_l) \in \mathfrak{Q}$, we put

$$\Omega(\bar{q}) := \bigsqcup_{s=1}^l \Omega(q_s),$$

where $\Omega(q) = \{\xi \in \mathbb{T}; \xi^q = \mathbf{1}\}$ is the group of all q -roots of $\mathbf{1}$, and the symbol \sqcup stands for a “disjoint union”.

The following notation will be useful: for any $\bar{q} = (q_1, \dots, q_l) \in \mathfrak{Q}$, we set

$$l(\bar{q}) = l \quad \text{and} \quad w(\bar{q}) = q_1 + \dots + q_l.$$

In particular, $\Omega(\bar{q})$ has cardinality $w(\bar{q})$.

We endow each finite group $\Omega(q)$ with its normalized Haar measure m_q (i.e. the normalized counting measure), and each $\Omega(\bar{q}) = \Omega(q_1, \dots, q_l)$ with the probability measure

$$m_{\bar{q}} = \frac{1}{l(\bar{q})} \sum_{s=1}^{l(\bar{q})} m_{q_s}.$$

Here, of course, $\Omega(q_s)$ is considered as a subset of $\Omega(\bar{q})$, so that the measures m_{q_s} are disjointly supported.

For any infinite sequence $\mathbf{q} = (\bar{q}_n)_{n \geq 1} \in \mathfrak{Q}^{\mathbb{N}}$, we put

$$\Omega(\mathbf{q}) := \prod_{n=1}^{\infty} \Omega(\bar{q}_n),$$

and we endow $\Omega(\mathbf{q})$ with the product measure

$$m_{\mathbf{q}} = \bigotimes_{n=1}^{\infty} m_{\bar{q}_n}.$$

Given $\mathbf{q} = (\bar{q}_n)_{n \geq 1} \in \mathfrak{Q}^{\mathbb{N}}$, it is not difficult to describe a “canonical” orthonormal basis for $L^2(\Omega(\mathbf{q}), m_{\mathbf{q}})$. However, we have to be careful with the notation.

For each sequence $\bar{q} = (q_1, \dots, q_l) \in \mathfrak{Q}$, let us denote by $\Gamma(\bar{q})$ the disjoint union $\bigsqcup_{s=1}^l \Gamma(q_s)$, where $\Gamma(q)$ is the dual group of $\Omega(q)$, i.e. $\Gamma(q) = \mathbb{Z}_q$. For any $\gamma_s \in$

$\Gamma(q_s) \subset \Gamma(\bar{q})$, define a function $e_{\gamma_s} : \Omega(\bar{q}) \rightarrow \mathbb{C}$ by

$$e_{\gamma_s}(\xi) = \begin{cases} \sqrt{l(\bar{q})} \langle \gamma_s, \xi \rangle & \text{if } \xi \in \Omega(q_s), \\ 0 & \text{otherwise.} \end{cases}$$

Since the characters of $\Omega(q)$ form an orthonormal basis of $L^2(\Omega(q), m_q)$ for every positive integer q , it is clear that the e_{γ_s} , $\gamma_s \in \Gamma(\bar{q})$ form an orthonormal basis of $L^2(\Omega(\bar{q}), m_{\bar{q}})$ for every $\bar{q} \in \mathfrak{Q}$.

Now, write each \bar{q}_n as $\bar{q}_n = (q_{n,1}, \dots, q_{n,l_n})$, and let us denote by $\Gamma(\mathbf{q})$ the set of all finite sequences of the form $\gamma = (\gamma_{s_1}, \dots, \gamma_{s_N})$ with $\gamma_{s_n} \in \Gamma(q_{n,s_n}) \subset \Gamma(\bar{q}_n)$ for all n and $\gamma_{s_N} \neq 0$. For any $\gamma = (\gamma_{s_1}, \dots, \gamma_{s_N}) \in \Gamma(\mathbf{q})$, we define $e_\gamma : \Omega(\mathbf{q}) \rightarrow \mathbb{C}$ as expected:

$$e_\gamma(\omega) = \prod_{n=1}^N e_{\gamma_{s_n}}(\omega_n).$$

In other words, $e_\gamma = e_{\gamma_{s_1}} \otimes \dots \otimes e_{\gamma_{s_N}}$. We also include the empty sequence \emptyset in $\Gamma(\mathbf{q})$ and we put $e_\emptyset = \mathbf{1}$. The following lemma is essentially obvious:

Lemma 5.1. *The family $(e_\gamma)_{\gamma \in \Gamma(\mathbf{q})}$ is an orthonormal basis of $L^2(\Omega(\mathbf{q}), m_{\mathbf{q}})$, for any $\mathbf{q} \in \mathfrak{Q}^{\mathbb{N}}$.*

We note that if $\mathbf{q} = (\bar{q}_n)_{n \geq 1} \in \mathfrak{Q}^{\mathbb{N}}$ and each sequence \bar{q}_n has length 1, i.e. $\bar{q}_n = (q_n)$ for some $q_n \geq 2$, then $(\Omega(\mathbf{q}), m_{\mathbf{q}})$ is just the compact group $\prod_{n \geq 1} \Omega(q_n)$ with its normalized Haar measure, and $\Gamma(\mathbf{q})$ “is” the character group of $\Omega(\mathbf{q})$. Moreover, if $E : \Omega(\mathbf{q}) \rightarrow X$ is a vector field on $(\Omega(\mathbf{q}), m_{\mathbf{q}})$ then $K_E(e_\gamma)$ is the Fourier coefficient $\widehat{E}(\gamma)$, for any $\gamma \in \Gamma(\mathbf{q})$. Accordingly, we shall use the following notation for an arbitrary $\mathbf{q} \in \mathfrak{Q}^{\mathbb{N}}$: given a vector field $E : \Omega(\mathbf{q}) \rightarrow X$ on $(\Omega(\mathbf{q}), m_{\mathbf{q}})$, we set

$$\widehat{E}(\gamma) = K_E(e_\gamma)$$

for every $\gamma \in \Gamma(\mathbf{q})$.

Next, we introduce a partially defined “metric” on every $\Omega(\mathbf{q})$, as follows. Write $\mathbf{q} = (\bar{q}_n)_{n \geq 1}$ and $\bar{q}_n = (q_{n,1}, \dots, q_{n,l_n})$. If $\omega = (\omega_n)$ and $\omega' = (\omega'_n)$ are distinct elements of $\Omega(\mathbf{q})$, let us denote by $n(\omega, \omega')$ the smallest n such that $\omega_n \neq \omega'_n$. If ω_n and ω'_n are in the same $\Omega(q_{n(\omega, \omega'), s})$, we denote this s by $s(\omega, \omega')$ and we put

$$d_{\mathbf{q}}(\omega, \omega') := \frac{1}{w(\bar{q}_1) \cdots w(\bar{q}_{n(\omega, \omega')-1})} \times \frac{1}{l_{n(\omega, \omega')}^{1/4} q_{n(\omega, \omega'), s(\omega, \omega')}}.$$

(The value of an empty product is declared to be 1). Otherwise, $d_{\mathbf{q}}(\omega, \omega')$ is not defined.

Definition 5.2. Let $\mathbf{q} \in \mathfrak{Q}^{\mathbb{N}}$. We shall say that a map $E : \Omega(\mathbf{q}) \rightarrow X$ is *super-Lipschitz* if

$$\|E(\omega) - E(\omega')\| \leq C d_{\mathbf{q}}(\omega, \omega')^2$$

for some finite constant C , whenever $d_{\mathbf{q}}(\omega, \omega')$ is defined.

The terminology is arguably not very convincing, since super-Lipschitz maps need not be continuous. Moreover, the definition of $d_{\mathbf{q}}$ may look rather special, mainly because of the strange term $l_{n(\omega, \omega')}^{1/4}$. We note, however, that when all sequences \bar{q}_n have length 1, say $\bar{q}_n = (q_n)$, then $d_{\mathbf{q}}$ is a quite natural true (ultrametric) distance on the Cantor-like group $\Omega(\mathbf{q}) = \prod_{n \geq 1} \Omega(q_n)$:

$$d_{\mathbf{q}}(\omega, \omega') = \frac{1}{q_1 \cdots q_{n(\omega, \omega')}} .$$

In this case, the terminology ‘‘super-Lipschitz’’ seems adequate (forgetting that we allowed arbitrary Hölder exponents $\alpha > 1$ in the weak mixing case). More importantly, the following lemma is exactly what is needed to carry out the first part of our program.

Lemma 5.3. *Let $\mathbf{q} \in \Omega^{\mathbb{N}}$. If $E : \Omega(\mathbf{q}) \rightarrow X$ is a super-Lipschitz vector field on $(\Omega(\mathbf{q}), m_{\mathbf{q}})$, then*

$$\sum_{\gamma \in \Gamma(\mathbf{q})} \|\widehat{E}(\gamma)\| < \infty .$$

Proof. Throughout the proof we put $\mathbf{q} = (\bar{q}_n)_{n \geq 1}$, and each \bar{q}_n is written as $\bar{q}_n = (q_{n,1}, \dots, q_{n,l_n})$.

For notational simplicity, we write Γ instead of $\Gamma(\mathbf{q})$. We denote by $|\gamma|$ the length of a sequence $\gamma \in \Gamma$. That is, $|\emptyset| = 0$ and $|\gamma| = N$ if $\gamma = (\gamma_{s_1}, \dots, \gamma_{s_N})$ with $\gamma_{s_n} \in \Gamma(q_{n,s_n})$ for all n and $\gamma_{N,s_N} \neq 0$. Then Γ is partitioned as

$$\Gamma = \bigcup_{N=0}^{\infty} \Gamma_N$$

where $\Gamma_N = \{\gamma \in \Gamma; |\gamma| = N\}$.

For any $\gamma = (\gamma_{s_1}, \dots, \gamma_{s_N}) \in \Gamma_N$, we put $s(\gamma) = s_N$. In other words, $s(\gamma)$ is the unique $s \in \{1, \dots, l_N\}$ such that the N -th coordinate of γ belongs to $\Omega(q_{N,s})$.

Fact. If $E : \Omega(\mathbf{q}) \rightarrow X$ is super-Lipschitz with constant C , then

$$\|\widehat{E}(\gamma)\| \leq \frac{C}{l_N} \sqrt{l_1 \cdots l_{N-1}} \prod_{n \leq N-1} w(\bar{q}_n)^{-2} \times \frac{1}{q_{N,s(\gamma)}^2}$$

for every $\gamma \in \Gamma_N$, $N \geq 1$.

Proof of Fact. Let us fix $\gamma = (\gamma_{s_1}, \dots, \gamma_{s_N}) \in \Gamma_N$, so that $s(\gamma) = s_N$. For notational simplicity (again), we put $q_{\gamma} = q_{N,s_N} = q_{N,s(\gamma)}$. For any $\xi \in \Omega(q_{\gamma})$ and $\omega \in \Omega(\mathbf{q})$ with $\omega_N \in \Omega(q_{\gamma})$, let us denote by $\xi\omega$ the element ω' of $\Omega(\mathbf{q})$ defined by $\omega'_n = \omega_n$ if

$n \neq N$ and $\omega'_N = \xi\omega_N$. Then

$$\begin{aligned} \widehat{E}(\gamma) &= \int_{\{\omega; \omega_N \in \Omega(q_\gamma)\}} e_\gamma(\omega) E(\omega) dm_{\mathbf{q}}(\omega) \\ &= \sum_{\xi \in \Omega(q_\gamma)} \int_{\{\omega; \omega_N = \xi\}} e_\gamma(\omega) E(\omega) dm_{\mathbf{q}}(\omega) \\ &= \sum_{\xi \in \Omega(q_\gamma)} \int_{\{\omega; \omega_N = \mathbf{1}_{\Omega(q_\gamma)}\}} e_\gamma(\xi\omega) E(\xi\omega) dm_{\mathbf{q}}(\omega) \\ &= \int_{\{\omega; \omega_N = \mathbf{1}_{\Omega(q_\gamma)}\}} e_\gamma(\omega) \times \left(\sum_{\xi \in \Omega(q_\gamma)} \langle \gamma_{s_N}, \xi \rangle E(\xi\omega) \right) dm_{\mathbf{q}}(\omega). \end{aligned}$$

Now, γ_{s_N} is assumed to be a non-trivial character of $\Omega(q_\gamma)$. So we have

$$\sum_{\xi \in \Omega(q_\gamma)} \langle \gamma_{s_N}, \xi \rangle = 0,$$

and it follows that

$$\widehat{E}(\gamma) = \int_{\{\omega; \omega_N = \mathbf{1}_{\Omega(q_\gamma)}\}} e_\gamma(\omega) \times \left[\sum_{\xi \in \Omega(q_\gamma)} \langle \gamma_{s_N}, \xi \rangle (E(\xi\omega) - E(\omega)) \right] dm_{\mathbf{q}}(\omega).$$

By the definition of a super-Lipschitz map, since $|e_\gamma(\omega)| \leq \sqrt{l_1 \cdots l_N}$ and since the set $\{\omega; \omega_N = \mathbf{1}_{\Omega(q_\gamma)}\}$ has $m_{\mathbf{q}}$ -measure $1/l_N q_\gamma$, we conclude that

$$\|\widehat{E}(\gamma)\| \leq \sqrt{l_1 \cdots l_N} \times \frac{1}{l_N} \times C \prod_{n \leq N-1} w(\bar{q}_n)^{-2} \times \frac{1}{l_N^{1/2} q_\gamma^2},$$

which is the required estimate. \square

It is now easy to conclude the proof of the lemma. For each $N \geq 1$ and every $s \in \{1, \dots, l_N\}$, the set $\Gamma_{N,s} = \{\gamma \in \Gamma_N; s(\gamma) = s\}$ has cardinality

$$|\Gamma_{N,s}| = \prod_{n \leq N-1} w(\bar{q}_n) \times q_{N,s}.$$

By the above fact and since $w(\bar{q}_n) = q_{n,1} + \cdots + q_{n,l_n} \geq 2l_n$ for all n , it follows that

$$\begin{aligned} \sum_{\gamma \in \Gamma_{N,s}} \|\widehat{E}(\gamma)\| &\leq \frac{C}{l_N} \sqrt{l_1 \cdots l_{N-1}} \prod_{n \leq N-1} w(\bar{q}_n)^{-1} \times \frac{1}{q_{N,s}} \\ &\leq \frac{C}{l_N} 2^{-N} \end{aligned}$$

for each $s \in \{1, \dots, l_N\}$. Hence, we get $\sum_{\gamma \in \Gamma_N} \|\widehat{E}(\gamma)\| \leq C 2^{-N}$ for every $N \geq 1$, and the result follows. \square

5.2. Construction of suitable \mathbb{T} -eigenfields. We now turn to the second part of our program, which is the most technical one. Our aim is to prove the following lemma.

Lemma 5.4. *Put $\mathbf{V} := \{(x, \lambda) \in X \times \mathbb{T}; x \neq 0 \text{ and } T(x) = \lambda x\}$, and let \mathbf{Z} be a closed subset of \mathbf{V} . Assume that for any (relatively) open set $O \subset \mathbf{V}$ such that $O \cap \mathbf{Z} \neq \emptyset$, the set $\{\lambda \in \mathbb{T}; \exists x : (x, \lambda) \in O\}$ is not \mathbf{S} -small. Then, for any (x_0, λ_0) in \mathbf{Z} and $\varepsilon > 0$, one can construct a \mathbb{T} -eigenfield (E, ϕ) for T on some $(\Omega(\mathbf{q}), m_{\mathbf{q}})$ such that*

- (1) $\|E(\omega) - x_0\| < \varepsilon$ for all $\omega \in \Omega(\mathbf{q})$;
- (2) $E : \Omega(\mathbf{q}) \rightarrow X$ is super-Lipschitz;
- (3) $\phi : \Omega(\mathbf{q}) \rightarrow \mathbb{T}$ is a homeomorphic embedding, and the image measure $\sigma = m_{\mathbf{q}} \circ \phi^{-1}$ is \mathbf{S} -continuous.

Proof. Throughout the proof, we denote (as usual) by $\mathcal{M}(\mathbb{T})$ the space of all complex measures on \mathbb{T} endowed with the total variation norm $\|\cdot\|$. We recall that $\mathcal{M}(\mathbb{T})$ is the dual space of $\mathcal{C}(\mathbb{T})$ (the space of all continuous complex-valued functions on \mathbb{T}), so we can use the w^* topology on $\mathcal{M}(\mathbb{T})$. Since our family \mathbf{S} is c_0 -like, we may fix a uniformly bounded sequence of w^* -continuous semi-norms $(\Phi_k)_{k \geq 0}$ on $\ell^\infty(\mathbb{Z}_+)$ such that

$$\mathbf{S} = \left\{ a \in \ell^\infty(\mathbb{Z}_+); \Phi_k(a) \xrightarrow{k \rightarrow \infty} 0 \right\}.$$

Without loss of generality, we may assume that $\Phi_k(a) \leq \|a\|_\infty$ for all k and every $a \in \ell^\infty(\mathbb{Z}_+)$. Moreover, upon replacing $\Phi_k(a)$ by $\max(|a_k|, \Phi_k(a))$, we may also assume that $\Phi_k(a) \geq |a_k|$ for every $a \in \ell^\infty(\mathbb{Z}_+)$.

Finally, to avoid notational heaviness, we denote by $\hat{\sigma}$ the *positive part* of the Fourier transform of a measure $\sigma \in \mathcal{M}(\mathbb{T})$, i.e. we write $\hat{\sigma}$ instead of $\hat{\sigma}|_{\mathbb{Z}_+}$ or $\mathcal{F}_+(\sigma)$. We then have

$$|\hat{\sigma}(k)| \leq \Phi_k(\hat{\sigma}) \leq \|\hat{\sigma}\|_\infty \leq \|\sigma\|$$

for all k and every $\sigma \in \mathcal{M}(\mathbb{T})$. In particular, if a bounded sequence $(\sigma_n) \subset \mathcal{M}(\mathbb{T})$ is such that the sequence $(\hat{\sigma}_n)$ is Cauchy with respect to each semi-norm Φ_k , then (σ_n) is w^* convergent in $\mathcal{M}(\mathbb{T})$.

Let us introduce some terminology. By an *admissible sequence of open sets in \mathbb{T}* , we shall mean a finite sequence of open sets $(V_i)_{i \in I} \subset \mathbb{T}$ such that the V_i have pairwise disjoint closures. An admissible sequence $(W_j)_{j \in J}$ is *finer* than an admissible sequence $(V_i)_{i \in I}$ if the following hold:

- for every $j \in J$, one can find $i \in I$ such that $\overline{W_j} \subset V_i$; then j is called a *successor* of i ;
- all $i \in I$ have the same number of successors $j \in J$.

In this situation, we write $i \prec j$ when $j \in J$ is a successor of $i \in I$. We define in the same way admissible sequences of open sets $(U_i)_{i \in I}$ in X and the corresponding refinement relation.

An *admissible pair* is a pair $(\sigma, (V_i)_{i \in I})$ where $(V_i)_{i \in I}$ is an admissible sequence of open sets in \mathbb{T} and σ is a positive \mathbf{S} -continuous measure such that

- $\text{supp}(\sigma) \subset \bigcup_{i \in I} V_i$;

- all open sets V_i have the same σ -measure.

An admissible pair $(\sigma, (V_i)_{i \in I})$ is *normalized* if σ is a probability measure.

Fact 1. Let $(\sigma, (V_i)_{i \in I})$ be a normalized admissible pair. Given $N \in \mathbb{N}$ and $\eta > 0$, one can find two admissible sequences of open sets $(V'_j)_{j \in J}$ and $(W_j)_{j \in J}$ (with the same index set J) and an \mathbf{S} -continuous probability measure ν such that

- $(V'_j)_{j \in J}$ is finer than $(V_i)_{i \in I}$ and $\text{diam}(V'_j) < \eta$ for all $j \in J$;
- $(\nu, (W_j)_{j \in J})$ is an admissible pair finer than $(\sigma, (V_i)_{i \in I})$;
- $\text{supp}(\nu) \subset \text{supp}(\sigma)$;
- $\text{supp}(\sigma) \cap V'_j \neq \emptyset$ for all $j \in J$;
- $\bigcup_j \overline{V'_j} \cap \bigcup_j \overline{W_j} = \emptyset$;
- $\|\nu - \sigma\| < \eta$;
- whenever σ' is a probability measure such that $(\sigma', (V'_j)_{j \in J})$ is an admissible pair, it follows that $\Phi_k(\widehat{\sigma}' - \widehat{\nu}) < \eta$ for all $k \leq N$.

Proof of Fact 1. Let η' and η'' be small positive numbers to be chosen later.

First, we note that since the measure σ is *continuous* by assumption on \mathbf{S} , one can partition $\text{supp}(\sigma) \cap V_i$, $i \in I$ into Borel sets $A_{i,1}, \dots, A_{i,K_i}$ with $\text{diam}(A_{i,s}) < \eta'$ and $\sigma(A_{i,1}) = \dots = \sigma(A_{i,K_i})$. This can be done as follows. Split $\text{supp}(\sigma) \cap V_i$ into finitely many Borel sets B_1, \dots, B_N with diameters less than $\eta'/2$ and positive σ -measure. Using the continuity of σ , choose a Borel set $B'_1 \subset B_1$ such that $\sigma(B'_1) > 0$ and $\sigma(B'_1)/\sigma(V_i)$ is a *rational* number. Then choose a Borel set B'_2 such that $B_1 \setminus B'_1 \subset B'_2 \subset (B_1 \setminus B'_1) \cup B_2$ and $\sigma(B'_2)/\sigma(V_i)$ is a positive rational number, and so on. This gives a partition of $\text{supp}(\sigma) \cap V_i$ into pairwise disjoint Borel sets B'_1, \dots, B'_N with diameter less than η' such that $\sigma(B'_k)/\sigma(V_i)$ is a positive rational number for each k , say $\sigma(B'_k) = \frac{p_k}{q} \sigma(V_i)$ where $p_k, q \in \mathbb{N}$ (the same q for all k). Finally, use again the continuity of σ to split each B'_k into p_k Borel sets $B''_{k,1}, \dots, B''_{k,p_k}$ with $\sigma(B''_{k,s}) = \frac{\sigma(V_i)}{q}$, and relabel the collection of all these sets $B''_{k,s}$ as $A_{i,1}, \dots, A_{i,K_i}$. That all the splittings can indeed be made follows from the (1-dimensional case of the) classical *Liapounoff convexity theorem*; see e.g. [51, Theorem 5.5].

Next, given any positive number m_i , one can partition further each $A_{i,s}$ into m_i Borel sets B with the same σ -measure. Taking $m_i := \prod_{j \neq i} K_j$, we then have the same number of Borel sets B inside each open set V_i . Thus, we may in fact assume from the beginning that we have the same number of sets $A_{i,s}$ inside each V_i . We denote this number by K , and we put $J := I \times \{1, \dots, K\}$. We note that $\sigma(A_{i,s}) = \frac{1}{|J|}$ for all $(i, s) \in J$.

Since the measure σ is regular and continuous, one can pick, for each $(i, s) \in J$, a compact set $C_{i,s} \subset A_{i,s}$ such that $0 < \sigma(A_{i,s} \setminus C_{i,s}) < \eta''$, and then a point $a_{i,s} \in (A_{i,s} \setminus C_{i,s}) \cap \text{supp}(\sigma)$. Then we may choose open sets $W_{i,s} \supset C_{i,s}$ with pairwise disjoint closures contained in V_i , and open sets $V'_{i,s} \ni a_{i,s}$ with pairwise disjoint closures contained in V_i and $\text{diam}(V'_{i,s}) < \eta'$, in such a way that $\bigcup_{j \in J} \overline{V'_j} \cap \bigcup_{j \in J} \overline{W_j} = \emptyset$.

If η'' is small enough, then the probability measure

$$\nu = \frac{1}{|J|} \sum_{(i,s) \in J} \frac{\sigma|_{C_{i,s}}}{\sigma(C_{i,s})}$$

satisfies $\|\sigma - \nu\| < \eta$. Moreover, $(\nu, (W_j)_{j \in J})$ is an admissible pair finer than $(\sigma, (V_i)_{i \in I})$.

Let us denote by ω_f the modulus of continuity of a function $f \in \mathcal{C}(\mathbb{T})$:

$$\omega_f(\delta) = \sup\{|f(u) - f(v)|; |u - v| < \delta\}.$$

Since the sets $C_{i,s}$, $(i, s) \in J$ form a partition of $\text{supp}(\nu)$ with $\nu(C_{i,s}) = \frac{1}{|J|}$, and since $|z - a_{i,s}| \leq \text{diam}(A_i, s) < \eta'$ for all $z \in C_{i,s}$, we have

$$\left| \int_{\mathbb{T}} f d\nu - \frac{1}{|J|} \sum_{(i,s) \in J} f(a_{i,s}) \right| \leq \omega_f(\eta')$$

for any $f \in \mathcal{C}(\mathbb{T})$. Similarly, if σ' is any probability measure with $\text{supp}(\sigma') \subset \bigcup_{j \in J} V'_j$ and $\sigma'(V'_j) = \frac{1}{|J|}$ for all $j \in J$, then

$$\left| \int_{\mathbb{T}} f d\sigma' - \frac{1}{|J|} \sum_{(i,s) \in J} f(a_{i,s}) \right| \leq \omega_f(\eta').$$

Hence, we see that σ' is close to ν in the w^* topology of $\mathcal{M}(\mathbb{T})$ if η' is sufficiently small. Since each semi-norm Φ_k is w^* -continuous and since the Fourier transformation $\mathcal{F}_+ : \mathcal{M}(\mathbb{T}) \rightarrow \ell^\infty(\mathbb{Z}_+)$ is (w^*, w^*) -continuous on bounded sets, it follows that if η' is small enough then, for any σ' such that $(\sigma', (V'_j)_{j \in J})$ is an admissible pair, we do have $\Phi_k(\hat{\sigma}' - \hat{\nu}) < \eta$ for all $k \leq N$. □

Remark. When considering two sequences of open sets $(V'_j)_{j \in J}$ and $(W_j)_{j \in J}$ both finer than a given sequence $(V_i)_{i \in I}$ and with the same index set J , we will always assume that the corresponding “extension” relations \prec on $I \times J$ are in fact the same.

At this point, we need to introduce some more terminology. By a *convenient triple*, we mean a triple $(\sigma, (U_i)_{i \in I}, (V_i)_{i \in I})$, where $(U_i) \subset X$ and $(V_i) \subset \mathbb{T}$ are admissible sequences of open sets (with the same index set I), and σ is a positive **S**-continuous measure such that the pair $(\sigma, (V_i)_{i \in I})$ is admissible and the following holds: for each $i \in I$ one can find a closed set $F_i \subset X$ such that

$$F_i \subset U_i \setminus \{0\} \text{ and } \text{supp}(\sigma) \subset \bigcup_{i \in I} \{\lambda \in V_i; \exists x \in F_i : (x, \lambda) \in \mathbf{Z}\}.$$

There is a natural notion of refinement for convenient triples, which is defined exactly as for admissible pairs. Moreover, we shall say that two convenient triples $(\sigma, (U_i), (V_i))$ and $(\sigma', (U_{i'}), (V_{i'}))$ are *disjoint* if $\bigcup_i \overline{U_i} \cap \bigcup_{i'} \overline{U_{i'}} = \emptyset$ and $\bigcup_i \overline{V_i} \cap \bigcup_{i'} \overline{V_{i'}} = \emptyset$.

The following fact is the key point in order to prove Lemma 5.4: it is the basic inductive step towards the construction of the measure space $(\Omega(\mathbf{q}), m_{\mathbf{q}})$ and the

map ϕ . The technical difficulty comes essentially from condition (c), which is here to ensure (3) in Lemma 5.4.

Fact 2. Let $(\sigma, (U_i)_{i \in I}, (V_i)_{i \in I})$ be a normalized convenient triple, and let $\eta > 0$. Then one can find a positive integer $l = l(\eta)$ and a finite sequence of pairwise disjoint normalized convenient triples $(\sigma'_1, (U'_j)_{j \in J_1}, (V'_j)_{j \in J_1}), \dots, (\sigma'_l, (U'_j)_{j \in J_l}, (V'_j)_{j \in J_l})$ finer than $(\sigma, (U_i), (V_i))$ such that

- (a) $\text{diam}(V'_j) < \eta$ for all $s \in \{1, \dots, l\}$ and every $j \in J_s$;
- (b) $\text{diam}(U'_j) < \frac{\eta}{l(\eta)^{1/2}|J_s|^2}$ for all s and every $j \in J_s$;
- (c) If we put

$$\sigma' := \frac{1}{l} \sum_{s=1}^l \sigma'_s$$

then $\sup_{k \geq 0} \Phi_k(\widehat{\sigma}' - \widehat{\sigma}) \leq \eta$.

Proof of Fact 2. Put $\nu_0 = \sigma = \sigma'_0$, $J_0 = I$ and $W_j = V_j$, $j \in J_0$. Let also η^* be a positive number to be chosen later but depending only on η , and let l be the smallest integer such that $l > 1/\eta^*$.

Since ν_0 is \mathbf{S} -continuous, one can find $N_0 \in \mathbb{N}$ such that $\Phi_k(\widehat{\nu}_0) < \eta^*$ for all $k > N_0$.

Applying Fact 1 to the pair $(\sigma, (V_i)_{i \in I}) = (\nu_0, (V_j)_{j \in J_0})$, we find an admissible sequence of open sets $(V'_j)_{j \in J_1}$ and a normalized admissible pair $(\nu_1, (W_j)_{j \in J_1})$ such that

- $(V'_j)_{j \in J_1}$ is finer than $(W_j)_{j \in J_0}$ and $\text{diam}(V'_j) < \eta^*$ for all $j \in J_1$;
- $(\nu_1, (W_j)_{j \in J_1})$ is finer than $(\nu_0, (W_j)_{j \in J_0})$;
- $\text{supp}(\nu_1) \subset \text{supp}(\nu_0)$;
- $\text{supp}(\nu_0) \cap V'_j \neq \emptyset$ for all $j \in J_1$;
- $\bigcup_{j \in J_1} \overline{V'_j} \cap \bigcup_{j \in J_1} \overline{W_j} = \emptyset$;
- $\|\nu_1 - \nu_0\| < \eta^*/l$;
- whenever σ' is a probability measure such that $(\sigma', (V'_j)_{j \in J_1})$ is an admissible pair, it follows that $\Phi_k(\widehat{\sigma}' - \widehat{\nu}_1) < \eta^*$ for all $k \leq N_0$.

Each $j \in J_1$ has a unique “predecessor” $i \in J_0$. In what follows, we denote this predecessor by j^- .

Since $V'_j \cap \text{supp}(\nu_0) \neq \emptyset$ and since $(\nu_0, (U_i)_{i \in J_0}, (V_i)_{i \in J_0})$ is a convenient triple, one can pick, for each $j \in J_1$, a point $\lambda_j \in V'_j \cap \text{supp}(\nu_0)$ and a point $x_j \in U'_{j^-}$ such that $(x_j, \lambda_j) \in \mathbf{Z}$. The points x_j are pairwise distinct because $x_j \neq 0$ and $T(x_j) = \lambda_j x_j$.

Using again the fact that $(\nu_0, (U_i)_{i \in J_0}, (V_i)_{i \in J_0})$ is a convenient triple, one can find closed sets $F_{i,0} \subset X$ with $F_{i,0} \subset U_i \setminus \{0\}$ such that

$$\text{supp}(\nu_0) \subset \bigcup_{i \in I} \{\lambda \in V_i; \exists x \in F_{i,0} : (x, \lambda) \in \mathbf{Z}\}.$$

Since \mathbf{Z} is closed in $(X \setminus \{0\}) \times \mathbb{T}$, the set

$$F_j = \{x \in F_{j^-,0}; \exists \lambda \in \text{supp}(\nu_1) \cap \overline{W_j} : (x, \lambda) \in \mathbf{Z}\}$$

is closed in X , for each $j \in J_1$. Moreover, the sets F_j are pairwise disjoint and we have $\bigcup_{j \in J_1} F_j \cap \{x_j; j \in J_1\} = \emptyset$, because $\bigcup_{j \in J_1} \overline{V'_j} \cap \bigcup_{j \in J_1} \overline{W_j} = \emptyset$. It follows that one can find open sets $U_j \subset X$, $j \in J_1$ with pairwise disjoint closures and open sets $U'_j \subset X$ with pairwise disjoint closures, such that

- $\text{diam}(U'_j) < \frac{\eta^*}{l^{1/2}|J_1|^2}$;
- $\overline{U_j} \subset U_{j^-}$ and $\overline{U'_j} \subset U_{j^-}$;
- $x_j \in U'_j$ and $F_j \subset U_j$;
- $\bigcup_{j \in J_1} \overline{U_j} \cap \bigcup_{j \in J_1} \overline{U'_j} = \emptyset$.

We note that $(\nu_1, (U_j)_{j \in J_1}, (W_j)_{j \in J_1})$ is a convenient triple, by the very definition of the closed sets F_j .

Now, we use the assumption on \mathbf{Z} : since $\mathbf{Z} \cap (U'_j \times V'_j) \neq \emptyset$, the set

$$A_j = \{\lambda \in V'_j; \exists x \in U'_j : (x, \lambda) \in \mathbf{Z}\}$$

is not \mathbf{S} -small, for any $j \in J_1$. Moreover, A_j is an analytic set because Z is a Borel subset of $X \times \mathbb{T}$; in particular, A_j is universally measurable (see [39]), and hence it contains a *compact* set which is not \mathbf{S} -small. Since the family of \mathbf{S} -continuous measures is hereditary with respect to absolute continuity, it follows that one can find, for each $j \in J_1$, an \mathbf{S} -continuous probability measure $\tilde{\sigma}_j$ such that

$$\text{supp}(\tilde{\sigma}_j) \subset \{\lambda \in V'_j; \exists x \in U'_j : (x, \lambda) \in \mathbf{Z}\}.$$

Moreover, since $U'_j \setminus \{0\}$ is an F_σ set in X , we may in fact assume that there is a closed set $F'_j \subset X$ with $F'_j \subset U'_j \setminus \{0\}$ such that

$$\text{supp}(\tilde{\sigma}_j) \subset \{\lambda \in V'_j; \exists x \in F'_j : (x, \lambda) \in \mathbf{Z}\}.$$

If we put

$$\sigma'_1 := \frac{1}{|J_1|} \sum_{j \in J_1} \tilde{\sigma}_j,$$

it follows that $(\sigma'_1, (U'_j)_{j \in J_1}, (V'_j)_{j \in J_1})$ is a convenient triple. In particular, the pair $(\sigma'_1, (V_j)_{j \in J_1})$ is admissible, so that $\Phi_k(\hat{\sigma}'_1 - \hat{\nu}_1) < \eta^*$ for all $k \leq N_0$.

Let us summarize what we have done up to now. Starting with the convenient triple $(\sigma, (U_i)_{i \in I}, (V_i)_{i \in I}) = (\nu_0, (U_j)_{j \in J_0}, (W_j)_{j \in J_0})$ and a positive integer N_0 such that $\Phi_k(\hat{\nu}_0) < \eta^*$ for all $k > N_0$, we have found two convenient triples $(\nu_1, (U_j)_{j \in J_1}, (W_j)_{j \in J_1})$ and $(\sigma'_1, (U'_j)_{j \in J_1}, (V'_j)_{j \in J_1})$ both finer than $(\sigma, (U_i), (V_i)) = (\nu_0, (U_j)_{j \in J_0}, (W_j)_{j \in J_0})$, such that

- $\bigcup_j \overline{V'_j} \cap \bigcup_j \overline{W_j} = \emptyset$;
- $\|\nu_1 - \nu_0\| < \eta^*/l$;
- $\Phi_k(\hat{\sigma}'_1 - \hat{\nu}_1) < \eta^*$ for all $k \leq N_0$.

Now, since ν_1 and σ'_1 are \mathbf{S} -continuous, we can choose $N_1 > N_0$ such that $\Phi_k(\hat{\nu}_1) < \eta^*$ and $\Phi_k(\hat{\sigma}'_1) < \eta^*$ for all $k > N_1$, and we repeat the whole procedure with $(\nu_1, (U_j)_{j \in J_1}, (W_j)_{j \in J_1})$ in place of $(\nu_0, (U_j)_{j \in J_0}, (W_j)_{j \in J_0})$. This produces two new normalized convenient triples $(\nu_2, (U_j)_{j \in J_2}, (W_j)_{j \in J_2})$ and $(\sigma'_2, (U'_j)_{j \in J_2}, (V'_j)_{j \in J_2})$.

Then we start again with ν_2 and a positive integer $N_2 > N_1$ witnessing that ν_2 and σ'_2 are \mathbf{S} -continuous, and so on.

After l steps, we will have constructed positive integers $N_0 < N_1 < \dots < N_l$ and, for each $s \in \{1, \dots, l\}$, two normalized convenient triples $(\nu_s, (U_j)_{j \in J_s}, (W_j)_{j \in J_s})$ and $(\sigma'_s, (U'_j)_{j \in J_s}, (V'_j)_{j \in J_s})$, such that the following properties hold:

- both triples $(\sigma'_s, (U'_j)_{j \in J_s}, (V'_j)_{j \in J_s})$ and $(\nu_s, (U_j)_{j \in J_s}, (W_j)_{i \in J_s})$ are finer than $(\nu_{s-1}, (U_j)_{j \in J_{s-1}}, (W_j)_{i \in J_{s-1}})$;
- $\text{diam}(U'_j) < \frac{\eta^*}{l^{1/2}|J_s|^2}$ and $\text{diam}(V'_j) < \eta^*$;
- $\bigcup_{j \in J_s} \overline{V'_j} \cap \bigcup_{j \in J_s} \overline{W_j} = \emptyset$;
- $\|\nu_l - \nu_{l-1}\| < \eta^*/l$;
- $\Phi_k(\hat{\sigma}'_s - \hat{\nu}_l) < \eta^*$ for all $k \leq N_{s-1}$;
- $\Phi_k(\hat{\nu}_s) < \eta^*$ and $\Phi_k(\hat{\sigma}'_s) < \eta^*$ for all $k > N_s$.

Then (a) and (b) hold, and we have to check (c). That is, we have to show that the measure

$$\sigma' = \frac{1}{l} \sum_{s=1}^l \sigma'_s,$$

satisfies $\Phi_k(\hat{\sigma}' - \hat{\sigma}) \leq \eta$ for all $k \geq 0$ if η^* is small enough (depending on η only) and $l > 1/\eta^*$.

First, we note that $\|\nu_s - \sigma\| = \|\nu_s - \nu_0\| < s\eta^*/l \leq \eta^*$ for every $s \in \{1, \dots, l\}$. Since $\Phi_k(\hat{\nu}_s - \hat{\sigma}) \leq \|\nu_s - \sigma\|$, it follows that

$$\Phi_k(\hat{\sigma}' - \hat{\sigma}) \leq \eta^* + \frac{1}{l} \sum_{s=1}^l \Phi_k(\hat{\sigma}'_s - \hat{\nu}_s)$$

for all $k \geq 0$.

If $k \leq N_0$ then $\Phi_k(\hat{\sigma}'_s - \hat{\nu}_s) < \varepsilon$ for every $s \in \{1, \dots, l\}$, and hence

$$\Phi_k(\hat{\sigma}' - \hat{\sigma}) \leq 2\eta^*.$$

If $k > N_0$, let us denote by $s(k)$ the largest $s \in \{0, \dots, l\}$ such that $k > N_s$. Then $\Phi_k(\hat{\sigma}'_s - \hat{\nu}_s) < \eta^*$ if $s > s(k) + 1$ because $k \leq N_{s(k)+1} \leq N_{s-1}$; and $\Phi_k(\hat{\sigma}'_s - \hat{\nu}_s) \leq \Phi_k(\hat{\sigma}'_s) + \Phi_k(\hat{\nu}_s) < 2\eta^*$ if $s \leq s(k)$, because $k > N_s$. So we get

$$\begin{aligned} \Phi_k(\hat{\sigma}' - \hat{\sigma}) &\leq \eta^* + \frac{1}{l} \left(\sum_{s \leq s(k)} 2\eta^* + \|\hat{\sigma}'_{s(k)+1} - \hat{\nu}_{s(k)+1}\|_\infty + \sum_{s > s(k)+1} \eta^* \right) \\ &\leq 4\eta^* + \frac{2}{l}. \end{aligned}$$

Taking $\eta^* = \eta/6$, this concludes the proof of Fact 2. \square

We can now start the actual proof of Lemma 5.4.

Recall that Ω is the set of all finite sequences of integers $\bar{q} = (q_1, \dots, q_l)$ with $q_s \geq 2$ for all s . We denote by $\Omega^{<\mathbb{N}}$ the set of all finite sequences $\mathbf{s} = (\bar{q}_1, \dots, \bar{q}_n)$ with $\bar{q}_j \in \Omega$, plus the empty sequence \emptyset . We denote by $|\mathbf{s}|$ the length of a sequence

$\mathbf{s} \in \Omega^{<\omega}$ (the empty sequence \emptyset has length 0), and by \prec the natural extension ordering on $\Omega^{<\omega}$.

If $\mathbf{s} = (\bar{q}_1, \dots, \bar{q}_n) \in \Omega^{<\mathbb{N}}$ we put

$$\Omega(\mathbf{s}) := \prod_{j=1}^n \Omega(\bar{q}_j),$$

and we denote by $m_{\mathbf{s}}$ the product measure $\otimes_{j=1}^n m_{\bar{q}_j}$. We also put $\Omega(\emptyset) = \{\emptyset\}$, and $m_{\emptyset} = \delta_{\emptyset}$.

Let us fix $(x_0, \lambda_0) \in \mathbf{Z}$ and $\varepsilon > 0$. Put $V_{\emptyset} := \mathbb{T}$, and choose an open set $U_{\emptyset} \subset X \setminus \{0\}$ such that $\text{diam}(U_{\emptyset}) < \varepsilon$ and $(x_0, \lambda_0) \in U_{\emptyset} \times V_{\emptyset}$. Let also pick an open set $U'_{\emptyset} \subset X$ such that $x_0 \in U'_{\emptyset}$ and $\overline{U'_{\emptyset}} \subset U_{\emptyset}$. By assumption, one can find an \mathbf{S} -continuous probability measure σ_0 such that $\text{supp}(\sigma_0) \subset \{\lambda \in V_{\emptyset}; \exists x \in U'_{\emptyset} : (x, \lambda) \in Z\}$. Then $(\sigma_0, U_{\emptyset}, V_{\emptyset})$ is a convenient triple.

We construct by induction a sequence $(\mathbf{s}_n)_{n \geq 0} \subset \Omega^{<\omega}$, a sequence of probability \mathbf{S} -continuous measures $(\sigma_n)_{n \geq 0}$ and, for each $n \geq 0$, two sequences of open sets $(U_{\xi})_{\xi \in \Omega(\mathbf{s}_n)} \subset X$ and $(V_{\xi})_{\xi \in \Omega(\mathbf{s}_n)} \subset \mathbb{T}$. If $n \geq 1$, we write $\mathbf{s}_n = (\bar{q}_1, \dots, \bar{q}_n)$ and $\bar{q}_j = (q_{j,1}, \dots, q_{j,l_j})$. If $\xi = (\xi_1, \dots, \xi_{n-1}) \in \Omega(\mathbf{s}_{n-1})$ and $\tau \in \Omega(\bar{q}_n)$, we denote by $\xi\tau$ the sequence $(\xi_1, \dots, \xi_{n-1}, \tau) \in \Omega(\mathbf{s}_n)$ (if $n = 1$ then $\xi\tau = \emptyset\tau = \tau$). Finally, we put $\varepsilon_0 = 1$ and

$$\varepsilon_n := \frac{1}{w(\bar{q}_1) \cdots w(\bar{q}_n)}$$

if $n \geq 1$. The following requirements have to be fulfilled for all $n \geq 0$.

- (i) $|\mathbf{s}_n| = n$, and $\mathbf{s}_{n-1} \prec \mathbf{s}_n$ if $n \geq 1$.
- (ii) If $n \geq 1$ and $\xi \in \Omega(\mathbf{s}_{n-1})$, then
 - $\overline{U_{\xi\tau}} \subset U_{\xi}$ and $\overline{V_{\xi\tau}} \subset V_{\xi}$ for every $\tau \in \Omega(\bar{q}_n)$;
 - $\overline{U_{\xi\tau}} \cap \overline{U_{\xi\tau'}} = \emptyset$ and $\overline{V_{\xi\tau}} \cap \overline{V_{\xi\tau'}} = \emptyset$ if $\tau \neq \tau'$;
 - $\text{diam}(U_{\xi\tau}) \leq \frac{1}{2} \text{diam}(U_{\xi})$ and $\text{diam}(V_{\xi\tau}) \leq \frac{1}{2} \text{diam}(V_{\xi})$.
- (iii) If $n \geq 1$ and $\xi \in \Omega(\mathbf{s}_{n-1})$, then

$$\text{diam}(U_{\xi\tau}) < \frac{\varepsilon_{n-1}}{l_n^{1/2} q_{n,s}^2}$$

for all $s \in \{1, \dots, l_n\}$ and every $\tau \in \Omega(q_{n,s})$.

- (iv) One can find closed sets $F_{\xi} \subset X$, $\xi \in \Omega(\mathbf{s}_n)$ with $F_{\xi} \subset U_{\xi} \setminus \{0\}$ and such that $\text{supp}(\sigma_n) \subset \bigcup_{\xi \in \Omega(\mathbf{s}_n)} \{\lambda \in V_{\xi}; \exists x \in F_{\xi} : (x, \lambda) \in \mathbf{Z}\}$. In particular:

$$\text{supp}(\sigma_n) \subset \bigcup_{\xi \in \Omega(\mathbf{s}_n)} \{\lambda \in V_{\xi}; \exists x \in U_{\xi} : (x, \lambda) \in \mathbf{Z}\}.$$

- (v) If $i \leq n$ and $\xi \in \Omega(\mathbf{s}_i)$, then $\sigma_n(V_{\xi}) = m_{\mathbf{s}_i}(\{\xi\})$.
- (vi) If $n \geq 1$ then $\sup_{k \geq 0} \Phi_k(\hat{\sigma}_n - \hat{\sigma}_{n-1}) < 2^{-n}$.

Since $(\sigma_0, U_{\emptyset}, V_{\emptyset})$ is a convenient triple, condition (iv) is satisfied for $n = 0$; and the other conditions are (trivially) satisfied as well.

Applying Fact 2 with the convenient triple $(\sigma_0, U_\emptyset, V_\emptyset)$ and a small enough $\eta > 0$, we get a positive integer l and a finite sequence of pairwise disjoint convenient triples $(\sigma'_1, (U'_j)_{j \in J_1}, (V'_j)_{j \in J_1}), \dots, (\sigma'_l, (U'_j)_{j \in J_l}, (V'_j)_{j \in J_l})$ finer than $(\sigma_0, U_\emptyset, V_\emptyset)$ such that

- $\text{diam}(U'_j) < \frac{1}{2} \text{diam}(U_\emptyset)$ and $\text{diam}(V'_j) < \frac{1}{2} \text{diam}(V_\emptyset)$ for all $s \in \{1, \dots, l\}$ and every $j \in J_s$;
- $\text{diam}(U'_j) < \frac{\varepsilon_0}{l^{1/2}|J_s|^2}$ for all s and every $j \in J_s$;
- $\sup_{k \geq 0} \Phi_k(\hat{\sigma}' - \hat{\sigma}_0) \leq 2^{-1}$, where $\sigma' = \frac{1}{l} \sum_{s=1}^l \sigma'_s$.

If we put $q_{1,s} := |J_s|$, $s \in \{1, \dots, l\}$ and $\bar{q}_1 = (q_{1,1}, \dots, q_{1,l})$, we may enumerate in the obvious way the open sets U'_j, V'_j as U_ξ, V_ξ , $\xi \in \Omega(\bar{q}_1)$. Then (i), ..., (vi) are clearly satisfied for $n = 1$ with $\sigma_1 = \sigma'$.

The general inductive step is very much the same. Assume that everything has been constructed up to some stage $n \geq 1$. Then, for every $\tilde{\xi} \in \Omega(\mathbf{s}_{n-1})$, the triple $\mathcal{T}_{\tilde{\xi}} = ((\sigma_n)_{|V_{\tilde{\xi}}}, (U_{\tilde{\xi}\tau})_{\tau \in \Omega(\bar{q}_n)}, (V_{\tilde{\xi}\tau})_{\tau \in \Omega(\bar{q}_n)})$ is a (non-normalized) convenient triple. Given $\eta > 0$, it is not hard to see (by examining the proof) that we can apply Fact 2 simultaneously to all triples $\mathcal{T}_{\tilde{\xi}}$, $\tilde{\xi} \in \Omega(\mathbf{s}_{n-1})$, with the same $l = l(\eta)$ and the same index set J . As above, we may take $J = \Omega(\bar{q}_n) \times \Omega(\bar{q})$, for some $\bar{q} = (q_1, \dots, q_l)$. This gives positive \mathbf{S} -continuous measures $\sigma_{\tilde{\xi}}$ and open sets $U_{\tilde{\xi}\tau'}$, $V_{\tilde{\xi}\tau'}$, $\tau' \in \Omega(\bar{q}_n) \times \Omega(\bar{q})$. Then, if η is small enough, conditions (i), ..., (vi) will be met at stage $n + 1$ with $\mathbf{s}_{n-1} = \mathbf{s}_n \bar{q}$, $\sigma_{n+1} = \sum_{\tilde{\xi}} \sigma_{\tilde{\xi}}$ and the open sets U_ξ, V_ξ for $\xi \in \Omega(\mathbf{s}_{n+1}) = \Omega(\mathbf{s}_{n-1}) \times \Omega(\bar{q}_n) \times \Omega(\bar{q})$.

Let us denote by \mathbf{q} the ‘‘limit’’ of the increasing sequence (\mathbf{s}_n) , i.e. the infinite sequence $(\bar{q}_n)_{n \geq 1} \in \Omega^{\mathbb{N}}$.

It follows from (ii) that for any $\omega = (\omega_n)_{n \geq 1} \in \Omega(\mathbf{q})$, the intersection $\bigcap_{n \geq 1} U_{\omega_n}$ is a single point $\{E(\omega)\}$, the intersection $\bigcap_{n \geq 1} V_{\omega_n}$ is a single point $\{\phi(\omega)\}$, and the maps $\phi : \Omega(\mathbf{q}) \rightarrow \mathbb{T}$ and $E : \Omega(\mathbf{q}) \rightarrow X$ are homeomorphic embeddings. Moreover, condition (iii) says exactly that E is super-Lipschitz. And since $x_0 \in U_\emptyset$ and $\text{diam}(U_\emptyset) < \varepsilon$, we have $\|E(\omega) - x_0\| < \varepsilon$ for every $\omega \in \Omega(\mathbf{q})$.

For each $\xi \in \bigcup_{n \geq 0} \Omega(\mathbf{s}_n)$, let us pick a point $\lambda_\xi \in V_\xi \cap \text{supp}(\sigma_n)$, where n is the length of ξ , i.e. $\xi \in \Omega(\mathbf{s}_n)$. By (iv), one can find $x_\xi \in U_\xi$ such that $(x_\xi, \lambda_\xi) \in \mathbf{Z}$. In particular, we have $T(x_\xi) = \lambda_\xi x_\xi$ for every $\xi \in \bigcup_{n \geq 0} \Omega(\mathbf{s}_n)$. Since $x_{\omega_n} \rightarrow E(\omega)$ and $\lambda_{\omega_n} \rightarrow \phi(\omega)$ as $n \rightarrow \infty$, it follows that $TE(\omega) = \phi(\omega) E(\omega)$ for every $\omega \in \Omega(\mathbf{q})$. In other words, (E, ϕ) is a \mathbb{T} -eigenfield for T .

By (vi), the sequence (σ_n) converges w^* to a probability \mathbf{S} -continuous measure σ . To conclude the proof, it remains to show that σ is the image measure of the measure $m_{\mathbf{q}}$ on $\Omega(\mathbf{q})$ under the embedding $\phi : \Omega(\mathbf{q}) \rightarrow \mathbb{T}$. That is, we want to check that

$$\int_{\Omega(\mathbf{q})} f \circ \phi(\omega) dm_{\mathbf{q}}(\omega) = \int_{\mathbb{T}} f d\sigma$$

for any $f \in \mathcal{C}(\mathbb{T})$. Let us fix f .

For each $\xi \in \bigcup_{n \geq 0} \Omega(\mathbf{s}_n)$, let us (again) pick a point $\lambda_\xi \in V_\xi$. Then $\lambda_{\omega|_n} \rightarrow \phi(\omega)$ as $n \rightarrow \infty$, for every $\omega \in \Omega(\mathbf{q})$. Setting $\Omega_\xi := \{\omega \in \Omega(\mathbf{q}); \xi \subset \omega\}$ and using Lebesgue's theorem, it follows that

$$\begin{aligned} \int_{\Omega(\mathbf{q})} f \circ \phi(\omega) dm_{\mathbf{q}}(\omega) &= \lim_{n \rightarrow \infty} \sum_{\xi \in \Omega(\mathbf{s}_n)} f(\lambda_\xi) \times m_{\mathbf{q}}(\Omega_\xi) \\ &= \lim_{n \rightarrow \infty} \sum_{\xi \in \Omega(\mathbf{s}_n)} f(\lambda_\xi) \times m_{\mathbf{s}_n}(\{\xi\}). \end{aligned}$$

On the other hand, by the definition of σ and since $\text{diam}(V_\xi) \rightarrow 0$ as $|\xi| \rightarrow \infty$ we also have

$$\begin{aligned} \int_{\mathbb{T}} f d\sigma &= \lim_{n \rightarrow \infty} \int_{\mathbb{T}} f d\sigma_n \\ &= \lim_{n \rightarrow \infty} \sum_{\xi \in \Omega(\mathbf{s}_n)} f(\lambda_\xi) \times \sigma_n(V_\xi). \end{aligned}$$

By condition (v), this concludes the proof. \square

5.3. The proof. Assume that the \mathbb{T} -eigenvectors are \mathbf{S} -perfectly spanning, and let us show that T is \mathbf{S} -mixing in the Gaussian sense. We recall that, since \mathbf{S} is c_0 -like, the \mathbb{T} -eigenvectors are in fact \mathbf{S} -perfectly spanning for analytic sets (see Remark 1 just after Corollary 2.9).

Let $\mathbf{V} := \{(x, \lambda) \in X \times \mathbb{T}; x \neq 0 \text{ and } T(x) = \lambda x\}$. Applying Lemma 3.6 to the family of \mathbf{S} -continuous measures, we get a closed set $\mathbf{Z} \subset \mathbf{V}$ such that

- for any $O \subset \mathbf{V}$ open such that $O \cap \mathbf{Z} \neq \emptyset$, the set $\{\lambda \in \mathbb{T}; \exists x : (x, \lambda) \in O\}$ is not \mathbf{S} -small;
- $\text{span}(\pi_1(\mathbf{Z}))$ is dense in X .

Using Lemma 5.3, Lemma 5.4 and proceeding exactly as in sub-section 4.4, we then find a sequence of \mathbb{T} -eigenfields (E_i, ϕ_i) defined on some $\Omega(\mathbf{q}_i)$, such that

- (1) each operator $K_{E_i} : L^2(\Omega(\mathbf{q}_i), m_{\mathbf{q}_i}) \rightarrow X$ is gamma-radonifying;
- (2) each E_i is continuous and $\overline{\text{span}}(\bigcup_{i \in \mathbb{N}} E_i[\Omega(\mathbf{q}_i)]) = X$;
- (3) each ϕ_i is a homeomorphic embedding, and $\sigma_i = m_{\mathbf{q}_i} \circ \phi_i^{-1}$ is \mathbf{S} -continuous.

Put $(\Omega_i, m_i) := (\Omega(\mathbf{q}_i), m_{\mathbf{q}_i})$ and let (Ω, m) be the disjoint union of the measure spaces (Ω_i, m_i) . Choose a sequence of small positive numbers $(\alpha_i)_{i \in \mathbb{N}}$, and define a \mathbb{T} -eigenfield (E, ϕ) on (Ω, m) as expected: $E(\omega_i) = \alpha_i E_i(\omega_i)$ and $\phi(\omega_i) = \phi_i(\omega_i)$ for each i and every $\omega_i \in \Omega_i$. By (1), the operator $K_E : L^2(\Omega, m) \rightarrow X$ is gamma-radonifying if the α_i are small enough. By (2) and since m has full support, the vector field E is m -spanning and hence the operator K_E has dense range. The intertwining equation $TK_E = K_E M_\phi$ holds by the definition of E . Finally, it follows from (3) that the measure $(f_i m_i) \circ \phi_i^{-1}$ is \mathbf{S} -continuous for each i and every $f_i \in L^1(\Omega_i, m_i)$. Hence, the measure $\sigma_f = (f m) \circ \phi^{-1}$ is \mathbf{S} -continuous for every $f \in L^1(\Omega, m)$. By Proposition 3.3, this shows that T is \mathbf{S} -mixing in the Gaussian sense.

5.4. Fréchet spaces. The above proof can be reproduced almost word for word in the Fréchet space setting. More precisely, the following changes should be made.

- Modify the definition of “super-Lipschitz”: a map $E : \Omega(\mathbf{q}) \rightarrow X$ is super-Lipschitz if $E : \Omega(\mathbf{q}) \rightarrow (X, \|\cdot\|)$ is super-Lipschitz for any continuous semi-norm $\|\cdot\|$ on X .
- In Lemma 5.3, add “for any continuous semi-norm $\|\cdot\|$ on X ”.
- In Lemma 5.4, fix a nondecreasing sequence of semi-norms $(\|\cdot\|_i)_{i \in \mathbb{N}}$ generating the topology of X , and put e.g.

$$\|x\| = \sum_{i \in \mathbb{N}} 2^{-i} \min(\|x\|_i, 1).$$

(Of course this is not even a semi-norm, but the notation is convenient anyway). Then perform exactly the same construction.

- Do the same when starting the proof of Theorem 2.6.

6. PROOF OF THE ABSTRACT RESULTS (2)

6.1. Proof of Theorem 2.8. Let the Banach space X have type 2. Let $T \in \mathfrak{L}(X)$ and assume that the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning for analytic sets. By Lemma 2.2 and Proposition 3.7, one can find a countable family of continuous \mathbb{T} -eigenvector fields $(E_i)_{i \in I}$ for T , where $E_i : \Lambda_i \rightarrow X$ is defined on some \mathbf{S} -perfect set $\Lambda_i \subset \mathbb{T}$, such that $\text{span}(\bigcup_{i \in I} E_i(\Lambda_i))$ is dense in X . By Lemma 2.3, each Λ_i is the support of some \mathbf{S} -continuous probability measure σ_i . If we put $\phi_i(\lambda) = \lambda$, this gives a continuous \mathbb{T} -eigenfield (E_i, ϕ_i) for T on the measure space (Λ_i, σ_i) such that the image measure $\sigma_i \circ \phi_i^{-1} = \sigma_i$ is \mathbf{S} -continuous; and by Proposition 3.1, the operator K_{E_i} is gamma-radonifying because X has type 2. So the proof can be completed exactly as in sub-section 5.3 above.

6.2. Proof of Proposition 2.7. Let the Banach space X have cotype 2, assume that $T \in \mathfrak{L}(X)$ is \mathbf{S} -mixing with respect to some Gaussian measure μ with full support, and let us show that the \mathbb{T} -eigenvectors of T are perfectly spanning for analytic sets.

We start with the following fact, which follows rather easily from Lemma 3.4 (1) (see the beginning of the proof of Theorem 5.46 in [9]).

Fact. One can find a gamma-radonifying operator $K : \mathcal{H} \rightarrow X$ such that $\mu = \mu_K$ and a unitary operator $M : \mathcal{H} \rightarrow \mathcal{H}$ such that $TK = KM$.

By the spectral theorem, we may assume that $\mathcal{H} = L^2(\Omega, m)$ for some measure space (Ω, m) and that M is a multiplication operator, $M = M_\phi$ for some measurable map $\phi : \Omega \rightarrow \mathbb{T}$. Then we use the cotype 2 assumption on X : by Proposition 3.1, the gamma-radonifying operator $K : L^2(\Omega, m) \rightarrow X$ is in fact given by a vector field; that is, $K = K_E$ for some vector field $E : (\Omega, m) \rightarrow X$.

Now, let $D \subset \mathbb{T}$ be an analytic \mathbf{S} -small set. By Remark 1 just after Lemma 3.4 and by Lemma 3.5 (with $\mathcal{H}_1 = \mathcal{H} \ominus \ker(K_E)$), we know that $\mathbf{1}_{\{\phi \in D\}} h \in \ker(K_E)$ for

any $h \in L^2(\Omega, m)$. In other words, we have

$$\int_{\{\phi \in D\}} h(\omega) E(\omega) dm(\omega) = 0$$

for every $h \in L^2(\Omega, m)$. It follows at once that E is almost everywhere 0 on the set $\{\phi \in D\}$; and since E is m -spanning (because $\mu = \mu_{K_E}$ has full support), this implies that the linear span of $\{E(\omega); \omega \in \Omega \setminus \phi^{-1}(D)\}$ is dense in X . By the very definition of a \mathbb{T} -eigenfield, it follows that the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in X , and we conclude that the \mathbb{T} -eigenvectors of T are \mathbf{S} -perfectly spanning.

7. CONCRETE EXAMPLES

Theorem 1.1 is a theoretical statement. However, it is extremely useful to give concrete examples of mixing operators, including new ones (e.g. the backward shift operators on c_0 , or the translation semigroups in sub-section 8.6 below). In this sub-section, we review several examples that were already known to be mixing in the Gaussian sense, either by an application of the results of [7], [8] or [9], or by using an ad-hoc argument. What we want to point out is that Theorem 1.1 now makes the proofs completely straightforward. In all cases, Theorem 1.1 is used through the following immediate consequence.

Proposition 7.1. *Let T be an operator acting on a complex separable Fréchet space X . Assume that one can find a \mathbb{T} -eigenfield (E, ϕ) for T on some topological measure space (Ω, m) such that the measure m has full support, E is continuous with $\overline{\text{span}} E(\Omega) = X$, and $(fm) \circ \phi^{-1}$ is a Rajchman measure for every $f \in L^1(m)$. Then T is strongly mixing in the Gaussian sense.*

Proof. If $D \subset \mathbb{T}$ is a Borel set of extended uniqueness, then $m(\phi^{-1}(D)) = 0$ because $(fm) \circ \phi^{-1}(D) = 0$ for every $f \in L^1(m)$. Since m has full support, it follows that $\Omega \setminus \phi^{-1}(D)$ is dense in Ω and hence that the linear span of $E(\Omega \setminus \phi^{-1}(D))$ is dense in X because E is assumed to be continuous. Since $E(\Omega \setminus \phi^{-1}(D)) \subset \bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$, this shows that the \mathbb{T} -eigenvectors of T are \mathcal{U}_0 -perfectly spanning. \square

Remark 1. When the measure m is finite, the third condition in Proposition 7.1 just means that $m \circ \phi^{-1}$ is a Rajchman measure.

Remark 2. The crucial difference with Proposition 3.3 is that the operator K_E is not assumed to be gamma-radonifying. So this is a much stronger result, but there is one drawback: it does not explain where the Gaussian measure comes from. On the other hand, what Proposition 3.3 says is that *if* we are lucky enough to have at hand a nice \mathbb{T} -eigenfield with a gamma-radonifying K_E , then the measure $\mu = \mu_{K_E}$ does the job.

7.1. Weighted shifts. Let $\mathbf{w} = (w_n)_{n \geq 1}$ be a bounded sequence of nonzero complex numbers, and let $B_{\mathbf{w}}$ be the associated *weighted backward shift* acting on $X_p = \ell^p(\mathbb{N})$, $1 \leq p < \infty$ or $X_{\infty} = c_0(\mathbb{N})$; that is,

$$B_{\mathbf{w}}(x_0, x_1, x_2, \dots) = (w_1 x_1, w_2 x_2, \dots).$$

Solving the equation $B_{\mathbf{w}}(x) = \lambda x$, it is easy to check that $B_{\mathbf{w}}$ has eigenvalues of modulus 1 if and only if

$$(7.1) \quad \text{the sequence } \left(\frac{1}{w_0 \cdots w_n} \right)_{n \geq 0} \text{ is in } X_p$$

(we have put $w_0 := 1$). In this case the formula

$$E(\lambda) := \sum_{n=0}^{\infty} \frac{\lambda^n}{w_0 \cdots w_n} e_n$$

defines a continuous \mathbb{T} -eigenvector field $E : \mathbb{T} \rightarrow X_p$ such that $\overline{\text{span}} E(\mathbb{T}) = X_p$. Hence $B_{\mathbf{w}}$ is strongly mixing in the Gaussian sense. This is known since [8] if $p < \infty$, but it appears to be new for weighted shifts on $c_0(\mathbb{N})$.

By Theorem 1.1, the converse is true if $p \leq 2$ (i.e. (7.1) is satisfied if $B_{\mathbf{w}}$ is strongly mixing in the Gaussian sense) since in this case X_p has cotype 2; but the case $p > 2$ is not covered by Theorem 1.1. However, it turns out that the converse does hold true for any $p < \infty$. In fact, (7.1) is satisfied as soon as the weighted shift $B_{\mathbf{w}}$ is frequently hypercyclic ([11]). As shown in [8], this breaks down completely when $p = \infty$: there is a frequently hypercyclic weighted shift $B_{\mathbf{w}}$ on $c_0(\mathbb{N})$ whose weight sequence satisfies $w_1 \cdots w_n = 1$ for infinitely many n . Such a weighted shift does not admit any (nontrivial) invariant Gaussian measure.

Let us also recall that, in contrast with the ergodic properties, the hypercyclicity of $B_{\mathbf{w}}$ does not depend on p : by a well known result of H. Salas ([54]), $B_{\mathbf{w}}$ is hypercyclic on X_p for any p if and only if $\sup_{n \geq 1} |w_1 \cdots w_n| = \infty$. Likewise, $B_{\mathbf{w}}$ is strongly mixing in the topological sense (on any X_p) iff $|w_1 \cdots w_n| \rightarrow \infty$. Hence, we see that strong mixing in the topological sense turns out to be equivalent to strong mixing in the Gaussian sense for weighted shifts on $c_0(\mathbb{N})$.

Finally, we would like to point out one quite curious fact. If $p < \infty$, the operator $K_E : L^2(\mathbb{T}) \rightarrow X_p$ turns out to be gamma-radonifying (see e.g. [9]) and hence there is a very natural explicit ergodic Gaussian measure for $B_{\mathbf{w}}$, namely the distribution of the random variable $\xi = \sum_0^{\infty} \frac{g_n}{w_0 \cdots w_n} e_n$. As shown in [8, Example 3.13], this need not be true when $p = \infty$, i.e. $X = c_0$; that is, the weight sequence can be chosen in such a way that K_E is not gamma-radonifying. So there is no “obvious” Gaussian measure in this case.

7.2. Kalisch operators. Let $X := \mathcal{C}_0([0, 2\pi]) = \{f \in \mathcal{C}([0, 2\pi]); f(0) = 0\}$ and let $V : L^2(0, 2\pi) \rightarrow X$ be the Volterra operator, $Vf(t) = \int_0^t f(s) ds$. There is a unique operator $T : X \rightarrow X$ such that $TV = VM_{\phi}$, where $M_{\phi} : L^2(0, 2\pi) \rightarrow L^2(0, 2\pi)$ is the multiplication operator associated with the function $\phi(t) = e^{it}$. The operator T is given by the formula

$$(7.2) \quad Tf(t) = \phi(t)f(t) - \int_0^t \phi'(s)f(s) ds.$$

It is easy to check that T has no eigenvalues. On the other hand, it can be shown that the operator V is gamma-radonifying (see e.g. [9]), and that the corresponding

Gaussian measure is in fact the classical *Wiener measure* on $\mathcal{C}_0([0, 2\pi])$ (see [44]). Hence (by Lemma 3.4), T is strongly mixing with respect to Wiener measure.

Consider now the operator T defined by formula (7.2), but acting on $L^2(0, 2\pi)$. It is straightforward to check that for any $t \in (0, 2\pi)$, the function $f_t = \mathbf{1}_{(0,t)}$ is an eigenvector for T with associated eigenvalue $\lambda = e^{it}$. Moreover the map $E : \mathbb{T} \setminus \{1\} \rightarrow L^2(0, 2\pi)$ defined by $E(e^{it}) = f_t$ is clearly continuous. Now, let Λ be an arbitrary compact subset of $\mathbb{T} \setminus \{1\}$. Let us denote by H_Λ the closed linear span of $E(\Lambda)$, and let T_Λ be the restriction of T to H_Λ . By a result of G. Kalisch ([37], see also [6, Lemma 2.12]), the point spectrum of T_Λ is exactly equal to Λ . By Corollary 1.2 and Theorem 1.1 (3), it follows that the operator T_Λ is weakly mixing in the Gaussian sense if and only if Λ is a perfect set, and strongly mixing iff Λ is \mathcal{U}_0 -perfect. Hence, any perfect \mathcal{U}_0 -set Λ gives rise to a very simple example of a weakly mixing transformation which is not strongly mixing. This could be of some interest since the classical concrete examples are arguably more complicated (see e.g. the one given in [50, section 4.5]).

Remark. Regarding the difference between weak and strong mixing, it is also worth pointing out that there exist Hilbert space operators which are weakly mixing in the Gaussian sense but not even strongly mixing in the topological sense. Indeed, in the beautiful paper [3], C. Badea and S. Grivaux are able to construct a weakly mixing operator (in the Gaussian sense) which is *partially power-bounded*, i.e. $\sup_{n \in I} \|T^n\| < \infty$ for some infinite set $I \subset \mathbb{N}$. This line of investigations was pursued even much further in [4] and [26].

7.3. Composition operators. Let $\alpha : \mathbb{D} \rightarrow \mathbb{D}$ be an automorphism of the unit disk \mathbb{D} without fixed points in \mathbb{D} , and let C_α be the associated composition operator acting on $X = H^p(\mathbb{D})$, $1 \leq p < \infty$:

$$C_\alpha(f) = f \circ \alpha.$$

As shown in [7], there is a natural continuous \mathbb{T} -eigenfield (E, ϕ) for C_α , defined on $\Omega = \mathbb{R}_+$ in the parabolic case and $\Omega = [0, 2\pi)$ in the hyperbolic case (endowed with Lebesgue measure m) such that $\overline{\text{span}} E(\Omega) = X$ and the image measure $m \circ \phi^{-1}$ is absolutely continuous with respect to Lebesgue measure on \mathbb{T} . By Proposition 7.1, it follows that C_α is strongly mixing in the Gaussian sense. This was shown in [8], with a longer proof because the authors had to play with the regularity of the vector field E and the geometry of the space X to show that K_E is gamma-radonifying; but the proof of [8] is also more informative since it provides an explicit Gaussian measure for C_α .

7.4. Operators with analytic \mathbb{T} -eigenvector fields. The following consequence of Proposition 7.1 is worth stating explicitly.

Lemma 7.2. *Let X be a complex separable Fréchet space, and let $T \in \mathfrak{L}(X)$. Assume that T is not a scalar multiple of the identity and that one can find a map $E : U \rightarrow X$ defined on some connected open set $U \subset \mathbb{C}$ such that E is holomorphic or “anti-holomorphic” (i.e. $E(\bar{s})$ is holomorphic), each $E(s)$ is an eigenvector for T ,*

the associated eigenvalue has modulus 1 for at least one $s \in U$ and $\overline{\text{span}} E(U) = X$. Then T is strongly mixing in the Gaussian sense.

Proof. Let us denote by $\phi(s)$ the eigenvalue associated with $E(s)$. Using the Hahn-Banach theorem, it is easily seen that $\phi : U \rightarrow \mathbb{C}$ is holomorphic or anti-holomorphic. Moreover, ϕ is non-constant since T is not scalar. By the inverse function theorem, it follows that $\phi(U) \cap \mathbb{T}$ contains a non trivial arc Λ such that the restriction of ϕ to $\Omega := \phi^{-1}(\Lambda)$ is a homeomorphism from Ω onto Λ . By the Hahn-Banach theorem and the identity principle for holomorphic (or anti-holomorphic) functions, the linear span of $E(\Omega)$ is dense in X . Hence, if we denote by m the image of the Lebesgue measure on Λ by ϕ^{-1} , the \mathbb{T} -eigenfield (E, ϕ) restricted to Ω satisfies the assumptions of Proposition 7.1.

One could also have used Theorem 1.1 directly. Indeed, if $D \subset \mathbb{T}$ is a Borel \mathcal{U}_0 -set, then $\phi^{-1}(\mathbb{T} \setminus D)$ is an uncountable Borel set (because $\phi(U)$ contains a non-trivial arc by the open mapping theorem), and as such it contains an uncountable compact set K . Then $\overline{\text{span}} E(K) = X$ by the identity principle, and since $E(K) \subset \bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$, it follows that the \mathbb{T} -eigenvectors of T are \mathcal{U}_0 -perfectly spanning. \square

This lemma may be applied, for example, in the following two cases.

- $T = M_\phi^*$, where M_ϕ is a (non-scalar) multiplication operator on some reproducing kernel Hilbert space of analytic functions on a connected open set $U \subset \mathbb{C}$, and $\phi(U) \cap \mathbb{T} \neq \emptyset$.
- T is a (non-scalar) operator on the space of entire functions $H(\mathbb{C})$ commuting with all translation operators.

In the first case one may take $E(s) := k_s$, the reproducing kernel at $s \in U$ (which depends anti-holomorphically on s and satisfies $M_\phi^*(k_s) = \overline{\phi(s)} k_s$). In the second case the assumptions of the lemma are satisfied with $E(s) = e_s$, where $e_s(z) = e^{sz}$ (denoting by τ_z the operator of translation by z , use the relation $\tau_z e_s = e_s(z) e_s$ to show that $T e_s = (T e_s(0)) \times e_s$).

Remark. Let Ω be a connected open set in \mathbb{C} . A Hilbert space operator $T \in \mathcal{L}(H)$ is a *Cowen-Douglas operator* with respect to Ω if $T - \omega$ is onto with $\dim \ker(T - \omega) < \infty$ for every $\omega \in \Omega$, and $\overline{\text{span}} \bigcup_{\omega \in \Omega} \ker(T - \omega) = H$. (This class of operators was introduced in [20]). It is shown in [57] that every such operator is unitarily equivalent to M_z^* acting on some reproducing kernel Hilbert space with underlying domain $U = \{\bar{\omega}; \omega \in \Omega\}$. Thus, we see that if $\Omega \cap \mathbb{T} \neq \emptyset$, then every Cowen-Douglas operator with respect to Ω is strongly mixing in the Gaussian sense.

8. MISCELLANEOUS REMARKS

8.1. The Frequent Hypercyclicity Criterion. An operator $T \in \mathcal{L}(X)$ satisfies the *Frequent Hypercyclicity Criterion* (see [7] and [15]) if one can find a set $\mathcal{D} \subset X$ with $\overline{\text{span}}(\mathcal{D}) = X$ and a map $S : \mathcal{D} \rightarrow \mathcal{D}$ such that

- (1) $TS(x) = x$ on \mathcal{D} ;

- (2) for every $x \in \mathcal{D}$, the series $\sum_{n \geq 0} T^n(x)$ and $\sum_{n \geq 0} S^n(x)$ are unconditionally convergent.

As the name indicates, an operator satisfying the Frequent Hypercyclicity Criterion is frequently hypercyclic. The original proof of this result is an explicit construction of a frequently hypercyclic vector, using a well-chosen sequence of subsets of \mathbb{N} with positive density. We know show that a straightforward application of Theorem 1.1 yields in fact a stronger conclusion.

Proposition 8.1. *If $T \in \mathfrak{L}(X)$ satisfies the Frequent Hypercyclicity Criterion, then T is strongly mixing in the Gaussian sense.*

Proof. For any $n \in \mathbb{Z}_+$, write $T^{-n} := S^n$. If $x \in \mathcal{D}$ then, by the unconditional convergence of the series in (2) above, the formula

$$E_x(\lambda) = \sum_{n \in \mathbb{Z}} \lambda^{-n} T^n(x)$$

makes sense for every $\lambda \in \mathbb{T}$ and defines a continuous map $E_x : \mathbb{T} \rightarrow X$. Moreover, it is immediately checked that $TE_x(\lambda) = \lambda E_x(\lambda)$, i.e. each E_x is a \mathbb{T} -eigenvectorfield for T . Finally, it follows easily from the Hahn-Banach theorem that the linear span of $\bigcup_{x \in \mathcal{D}} E_x(\mathbb{T})$ is dense in X . Hence, we may apply Corollary 1.2. \square

Remark 1. A similar result was obtained independently by M. Murillo-Arcilla and A. Peris [45]. Namely, it is shown in [45] that if T satisfies the Frequent Hypercyclicity Criterion, then one can find a Borel probability measure μ on X with full support such that T is mixing with respect to μ . Interestingly enough, the measure μ is *not* Gaussian. In [46], the same authors also obtained the existence of mixing measures with full support for one-parameter semigroups satisfying the Frequent Hypercyclicity Criterion.

Remark 2. Consider the derivation operator D on the space of entire functions $H(\mathbb{C})$. There is a canonical right inverse $S : H(\mathbb{C}) \rightarrow H(\mathbb{C})$, namely $Sf(z) = \int_{[0,z]} f(\xi) d\xi$. Putting $\mathcal{D} = \{S^n \mathbf{1}; n \geq 0\} = \{\frac{z^n}{n!}; n \geq 0\}$, assumptions (1), (2) above are satisfied. With the notations of the above proof we have $E_1(\lambda) = \sum_{n=0}^{\infty} \lambda^{-n} \frac{z^n}{n!}$, and the linear span of $E_1(\mathbb{T})$ is already dense in \mathbb{T} . Moreover, the operator $K_{E_1} : L^2(\mathbb{T}) \rightarrow H(\mathbb{C})$ is gamma-radonifying. Denoting by $(e_n)_{n \in \mathbb{Z}}$ the trigonometric basis of $L^2(\mathbb{T})$, we have $K_{E_1} e_n = \widehat{E_1}(n) = 0$ if $n > 0$ and $\frac{z^n}{n!}$ if $n \geq 0$; so the corresponding Gaussian measure is $\mu \sim \sum_0^{\infty} g_n \frac{z^n}{n!}$. Thus, we arrive at the following conclusion: *the random entire function $f = \sum_0^{\infty} g_n \frac{z^n}{n!}$ is almost surely frequently hypercyclic for the derivation operator D .* In fact, this is true for essentially any well-defined random entire function of the form $\sum_0^{\infty} \xi_n \frac{z^n}{n!}$, where (ξ_n) is an identically distributed sequence of symmetric random variables satisfying some simple nontriviality condition. This was shown recently by M. Nikula [?].

8.2. Non Gaussian measures. It is natural to ask whether one gets a more general notion of mixing for an operator $T \in \mathfrak{L}(X)$ by requiring only that T should be mixing with respect to *some* probability measure μ on X with full support. The following remark (essentially contained in [52], and also in [7]) provides a partial answer.

Remark. Let X be a separable Banach space, and assume that X has *type 2*. Let also μ be a centred Borel probability measure on X such that $\int_X \|x\|^2 d\mu(x) < \infty$. Then there is a unique Gaussian measure ν on X such that

$$\|x^*\|_{L^2(\nu)} = \|x^*\|_{L^2(\mu)}$$

for every $x^* \in X^*$. Moreover,

- (1) if μ has full support then so does ν ;
- (2) if μ is T -invariant for some $T \in \mathfrak{L}(X)$, then so is ν ;
- (3) if $T \in \mathfrak{L}(X)$ is weakly mixing or strongly mixing with respect to μ , then the same is true with respect to ν .

Proof. Put $\mathcal{H} := L^2(\mu)$. By assumption on μ , there is a well defined (conjugate-linear) “inclusion” operator $J : X^* \rightarrow \mathcal{H}$, namely $J(x^*) = \overline{x^*}$ considered as an element of $L^2(\mu)$. Moreover, it follows from Lebesgue’s theorem and the w^* -metrizable of B_{X^*} that J is (w^*, w^*) -continuous on B_{X^*} . Hence, J is the adjoint of a bounded operator $K : \mathcal{H} \rightarrow X$. By definition, this means that $K^*x^* = \overline{x^*}$ considered as an element of $L^2(\mu)$, so we have in particular

$$\|K^*x^*\|_{\mathcal{H}} = \|x^*\|_{L^2(\mu)}.$$

It is fairly easy to show that the “inclusion” operator $K^* = J : X^* \rightarrow L^2(\mu)$ is *absolutely 2-summing*, i.e.

$$\sum_{i \in I} \|J\phi_i\|^2 \leq C \sup_{\phi^* \in B_{E^*}} \sum_{i \in I} |\langle \phi^*, \phi_i \rangle|^2$$

for some constant $C < \infty$ and every finite sequence $(\phi_i)_{i \in I} \subset E = X^*$ (see e.g. [2] for more on 2-summing operators). Since X has type 2, it follows that K is gamma-radonifying (see [17], or [9, Proposition 5.19]). If we denote by $\nu = \mu_K$ the associated Gaussian measure, then $\|x^*\|_{L^2(\nu)} = \|x^*\|_{L^2(\mu)}$ for every $x^* \in X^*$ by the very definition of ν . Moreover, if ν' is another Gaussian measure with the same properties, then ν and ν' have the same Fourier transform and hence $\nu = \nu'$.

To prove (1), assume that μ has full support. Then the operator $K^* = J : X^* \rightarrow \mathcal{H}$ is one-to-one because a continuous function on X is 0 in $L^2(\mu)$ if and only if it is identically 0. It follows that K has dense range, and hence that $\nu = \mu_K$ has full support.

To prove (2), assume that μ is T -invariant. Then $\|T^*x^*\|_{L^2(\mu)} = \|x^*\|_{L^2(\mu)}$ and hence $\|K^*(T^*x^*)\|_{L^2(\nu)} = \|K^*(x^*)\|_{L^2(\nu)}$ for every $x^* \in X^*$. By the proof of Lemma 3.4, this shows that ν is T -invariant

Finally, (3) follows from the following observations : (i) weak mixing or strong mixing of T with respect to μ is characterized by a certain behaviour (B) of the sequence $(\langle f \circ T^n, g \rangle_{L^2(\mu)})_{n \geq 0}$, for any $f, g \in L^2_0(\mu)$; (ii) when specialized to linear functionals, this gives that the sequence $(\langle T^{*n}x^*, y^* \rangle_{L^2(\mu)})$ satisfies (B) for any $x, y^* \in X^*$; (iii) by the definition of ν , this means that $(\langle T^{*n}x^*, y^* \rangle_{L^2(\nu)})$ satisfies (B) for any x^*, y^* ; (iv) since ν is a Gaussian measure, this is enough to ensure weak mixing or strong mixing of T with respect to ν , by Rudnicki [52] or Bayart-Grivaux [7]. \square

When X is a Hilbert space, it follows from this remark (and from Theorem 1.1) that an operator $T \in \mathfrak{L}(X)$ is e.g. strongly mixing with respect to some centred probability measure μ on X with full support such that $\int_X \|x\|^2 d\mu(x) < \infty$ if and only if the T -eigenvectors of T are \mathcal{U}_0 -perfectly spanning, in which case the measure μ can be assumed to be Gaussian. It would be interesting to know if this remains true without any a priori assumption on the measure μ .

8.3. A characterization of \mathcal{U}_0 -sets. Our results easily yield the following curious characterization of sets of extended uniqueness.

Remark. Let D be an analytic subset of \mathbb{T} . Then D is a set of extended uniqueness if and only if the following holds: for every Hilbert space operator T which is strongly mixing in the Gaussian sense, the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in the underlying Hilbert space.

Proof. The “only if” part follows immediately from Proposition 2.7. Conversely, assume that $D \notin \mathcal{U}_0$. Then, since D is universally measurable, it contains a compact set Λ which is the support of some Rajchman probability measure, i.e. Λ is \mathcal{U}_0 -perfect. Let $T_\Lambda : H_\Lambda \rightarrow H_\Lambda$ be the associated Kalisch operator (see sub-section 7.2). Then T_Λ is strongly mixing in the Gaussian sense, but $\text{span} \left(\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda) \right)$ is certainly not dense in H since it is $\{0\}$ (recall that $\sigma_p(T) = \Lambda$ and $\Lambda \subset D$). \square

8.4. A “non-ergodicity index”. Let us say that a Gaussian measure μ is *non-trivial* if $\mu \neq \delta_0$. From Theorem 1.1 (and since ergodicity and weak mixing in the Gaussian sense are equivalent) we immediately get the following result.

Corollary 8.2. *For a linear operator T acting on a Banach space X with cotype 2, the following are equivalent:*

- (a) T admits a nontrivial ergodic Gaussian measure;
- (b) there exists a closed, T -invariant subspace $Z \neq \{0\}$ such that

$$\overline{\text{span}} \bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T|_Z - \lambda) = Z$$

for every countable set $D \subset \mathbb{T}$.

In this case, T admits an ergodic Gaussian measure with support Z , for any such subspace Z .

Proof. If T admits an ergodic Gaussian measure $\mu \neq \delta_0$, then $Z := \text{supp}(\mu)$ is a non-zero T -invariant subspace, and Z satisfies (b) by Theorem 1.1 (3). The converse follows from Theorem 1.1 (1). \square

Loosely speaking, this result is a kind of “perfect set theorem” for ergodicity. This can be made precise as follows. Let $T \in \mathfrak{L}(X)$ (where X is a Banach space with cotype 2), and consider the following “derivation” on closed, T -invariant subspaces of X : for any such subspace E , set

$$\mathcal{D}_T(E) := \bigcap_D \overline{\text{span}} \bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T|_E - \lambda),$$

where the intersection ranges over all countable sets $D \subset \mathbb{T}$. By transfinite induction, one defines the iterates $\mathcal{D}_T^\alpha(X)$ for every ordinal α , in the obvious way:

$$\begin{aligned}\mathcal{D}_T^{\alpha+1}(X) &= \mathcal{D}_T[\mathcal{D}_T^\alpha(X)], \\ \mathcal{D}_T^\lambda(X) &= \bigcap_{\alpha < \lambda} \mathcal{D}_T^\alpha(X) \quad (\lambda \text{ limit}).\end{aligned}$$

Since X is a Polish space, the process must stabilize at some countable ordinal $\alpha(T)$ and hence $\mathcal{D}_T^\infty(X) := \bigcap_\alpha \mathcal{D}_T^\alpha(X)$ is well-defined and is a fixed point of \mathcal{D}_T (in fact, the largest fixed point). Then, Corollary 8.2 says that T admits a nontrivial ergodic Gaussian measure if and only if $\mathcal{D}_T^\infty(X) \neq \{0\}$, in which case one can find an ergodic measure with support $\mathcal{D}_T^\infty(X)$. The subspace $\mathcal{D}_T^\infty(X)$ is the “perfect kernel” associated with the derivation \mathcal{D}_T , a canonical witness of the ergodicity of T .

Let us say that T is *totally non-ergodic* if it does not admit any nontrivial ergodic Gaussian measure. Then the ordinal $\alpha(T)$ may be called the “non-ergodicity index” of T . It is quite natural to wonder whether this index can be arbitrarily large: given any countable ordinal α , is it possible to construct a totally non-ergodic operator T with $\alpha(T) > \alpha$?

8.5. The scope of the abstract results. Some comments are in order regarding the assumptions made on the family \mathbf{S} in Theorems 2.6 and 2.8.

The main trouble with Theorem 2.6 is that we have no idea of how to prove it without assuming that the family \mathbf{S} is c_0 -like. Perhaps unexpectedly, what makes the definition of a c_0 -like family very restrictive is the uniform boundedness assumption of the sequence of semi-norms (Φ_n) . For example, any growth condition of the form

$$a_n = o(\varepsilon_n),$$

where $\bar{\varepsilon} = (\varepsilon_n)$ is a sequence of positive numbers tending to 0 and satisfying $\varepsilon_{n \pm k} \leq C_k \varepsilon_n$, defines a translation-invariant ideal $\mathbf{S}_{\bar{\varepsilon}} \subset \ell^\infty(\mathbb{Z}_+)$ which has the correct form *except* for this uniform boundedness condition (just put $\Phi_n(a) = \frac{1}{\varepsilon_n} |a_n|$). In fact, in this case one cannot hope for positive results of any kind: as observed by V. Devinck ([25]) it follows from a result of C. Badea and V. Müller ([5]) that $\mathbf{S}_{\bar{\varepsilon}}$ -mixing operators just do not exist at all (at least on a Hilbert space).

In the same spirit, it can be shown that it is impossible to find any c_0 -like description for $\mathcal{B} = L^1(m)$, the family of all measures absolutely continuous with respect to Lebesgue measure on \mathbb{T} (see [43]).

As a more extreme example, the well studied notion of *mild mixing* (see e.g. [1]) does not fit at all into the framework. Indeed, in this case the relevant family of measures \mathcal{B} is the following: a measure σ is in \mathcal{B} iff it annihilates every *weak Dirichlet set* (a Borel set $D \subset \mathbb{T}$ is weak Dirichlet if, for any measure μ supported on D , one can find an increasing sequence of integer (n_k) such that $z^{n_k} \rightarrow \mathbf{1}$ in $L^2(\mu)$). By a result of S. Kahane ([36]), the family \mathcal{B} is extremely complicated, namely *non Borel* in $(\mathcal{M}(\mathbb{T}), w^*)$. So there is absolutely no hope of finding a c_0 -like description for \mathcal{B} .

Now, if one is interested in Hilbert spaces only, Theorem 2.8 is arguably rather general since the c_0 -like property is not required. However, that the family \mathbf{S} should

be *norm-closed* in $\ell^\infty(\mathbb{Z}_+)$ is already quite a restrictive condition: indeed, this implies that the strongest mixing property that can be reached is, precisely, strong mixing. In particular, Theorem 2.8 cannot be applied to any “summability” condition on the Fourier coefficients. On the other hand, the following example can be handled: let \mathcal{F} be any translation-invariant filter on \mathbb{Z}_+ , and take as \mathbf{S} the family of all sequences $(a_n) \in \ell^\infty(\mathbb{Z}_+)$ tending to 0 along \mathcal{F} .

Perhaps more importantly, there are quite natural (norm-closed) families \mathbf{S} which are not algebras. The most irritating example is the ergodic case. As already observed a set $D \subset \mathbb{T}$ is \mathbf{S}_{erg} -small if and only if $D \subset \{1\}$. Hence, if Theorem 2.8 could be applied in this case, the result would read as follows: *if $T \in \mathfrak{L}(X)$ and if $\text{span}\left(\bigcup_{\lambda \in \mathbb{T} \setminus \{1\}} \ker(T - \lambda)\right)$ is dense in X , then T is ergodic in the Gaussian sense.* But this is clearly not true since for example $T = -id$ satisfies the assumption and is not even hypercyclic.

In spite of that, it might happen (as already suggested in [27]) that an operator $T \in \mathfrak{L}(X)$ is ergodic in the Gaussian sense as soon as *it is hypercyclic* and the \mathbb{T} -eigenvectors span X ; but the proof of such a result would require at least one new idea. This is, of course, related to the following well known open problem: is it true that every *chaotic* operator (i.e. a hypercyclic operator with a dense set of periodic points) is frequently hypercyclic?

8.6. Semigroups. The proof of Theorem 1.1 can be easily adapted to get analogous results in the continuous case, i.e. for one-parameter semigroups of operators $(T_t)_{t \geq 0}$. Indeed, the mixing properties make perfect sense in the continuous case, and the sets of extended uniqueness are well defined on \mathbb{R} just like on any locally compact abelian group. The only “difference” is that, according to the (point-)spectral mapping theorem for C_0 -semigroups, the unimodular eigenvalues of the single operator T from the discrete case should be replaced with the purely imaginary eigenvalues of the semigroup generator in the continuous case. Hence, the continuous analogue of Theorem 1.1 reads as follows.

Theorem 8.3. *Let X be a separable Banach space and let $\mathcal{T} = (T_t)_{t \geq 0}$ be a C_0 -semigroup in $\mathfrak{L}(X)$ with infinitesimal generator A .*

- (1) *If the linear span of $\bigcup_{\theta \in \mathbb{R} \setminus D} \ker(A - i\theta)$ is dense in X for any countable set $D \subset \mathbb{R}$, then \mathcal{T} is weakly mixing in the Gaussian sense.*
- (2) *If the linear span of $\bigcup_{\theta \in \mathbb{R} \setminus D} \ker(A - i\theta)$ is dense in X for any \mathcal{U}_0 -set $D \subset \mathbb{R}$, then \mathcal{T} is strongly mixing in the Gaussian sense.*

Since the proof would essentially be a matter of changing the notation, we shall not give any detail. We note, however, that the semigroup case is obviously of some interest in view of its connections with partial differential equations. See e.g. [16], [41] or [53] for more on these matters.

In another direction, a perhaps ambitious program would be to consider linear representations of more general semigroups; in other words, to establish results like Theorem 1.1 for semigroups of operators $(T_\gamma)_{\gamma \in \Gamma}$ with Γ no longer equal to \mathbb{N} or \mathbb{R}_+ . There is no difficulty in defining ergodicity or strong mixing in this setting, and the

spectral approach makes sense e.g. if Γ is a locally compact abelian group. However, it is not quite clear what the correct “perfect spanning” property should be.

8.6.1. *Translation semigroups.* To illustrate Theorem 8.3, let us give a new and very simple example of a strongly mixing C_0 -semigroup, that cannot be reached by applying the results of [7], [8] or [9].

Let $\rho : \mathbb{R}_+ \rightarrow (0, \infty)$ be a locally bounded positive function on \mathbb{R}_+ , and let

$$\mathcal{C}_0(\mathbb{R}_+, \rho) := \left\{ f \in \mathcal{C}(\mathbb{R}_+); \lim_{x \rightarrow +\infty} f(x)\rho(x) = 0 \right\}$$

endowed with its natural norm, $\|f\| = \|f\rho\|_\infty$. Moreover, assume that ρ is an *admissible weight* in the sense of [24], which means that

$$C(t) := \sup_{x \in \mathbb{R}_+} \frac{\rho(x)}{\rho(x+t)} < \infty$$

for any $t \geq 0$ and $C(t)$ is locally bounded on \mathbb{R}_+ . Then the *translation semigroup* $\mathcal{T} = (T_t)_{t \geq 0}$ defined by

$$T_t f(x) = f(x+t)$$

is a C_0 -semigroup on $\mathcal{C}_0(\mathbb{R}_+, \rho)$.

It is proved in [12] that \mathcal{T} is strongly mixing in the topological sense if and only if $\rho(x) \xrightarrow{x \rightarrow \infty} 0$ as $x \rightarrow \infty$. We now show that this is in fact equivalent to strong mixing in the Gaussian sense.

Proposition 8.4. *The translation semigroup \mathcal{T} is strongly mixing in the Gaussian sense on $\mathcal{C}_0(\mathbb{R}_+, \rho)$ if and only if $\rho(x) \rightarrow 0$ as $x \rightarrow +\infty$.*

Proof. Assume that $\rho(x) \rightarrow 0$ as $x \rightarrow +\infty$. The infinitesimal generator of \mathcal{T} is the derivation operator (denoted by A), whose domain includes all \mathcal{C}^1 functions on \mathbb{R}_+ with a bounded and uniformly continuous derivative. For any $\theta \in \mathbb{R}$, the function $e_\theta(x) = e^{i\theta x}$ is in $\ker(A - i\theta)$, and the map $\theta \mapsto e_\theta$ is clearly continuous from \mathbb{R} into $\mathcal{C}_0(\mathbb{R}_+, \rho)$. Moreover, it follows easily from (the Hahn-Banach theorem and) the injectivity of the Fourier transformation that the linear span of the functions e_θ is dense in $\mathcal{C}_0(\mathbb{R}_+, \rho)$. By Theorem 8.3, we conclude that \mathcal{T} is strongly-mixing in the Gaussian sense. \square

Remark. One may also consider the weighted L^p spaces $L^p(\mathbb{R}_+, \rho)$ ($1 \leq p < \infty$) defined by the condition

$$\int_0^\infty |f(x)|^p \rho(x) dx < \infty.$$

With exactly the same proof as above, one gets that the translation semigroup is strongly mixing in the Gaussian sense on $L^p(\mathbb{R}_+, \rho)$ as soon as

$$\int_0^\infty \rho(x) dx < \infty.$$

8.7. The size of the set of hypercyclic vectors. It is well known that if T is a hypercyclic operator acting on a separable Fréchet space X , then $HC(T)$ (the set of hypercyclic vectors for T) is a dense G_δ subset of X . Moreover, as observed in the introduction, if T happens to be ergodic with respect to some probability measure μ with full support, then μ -almost every $x \in X$ is a hypercyclic vector for T . Thus, $HC(T)$ is large both in the Baire category sense and in a measure-theoretic sense.

Now, there are many other natural notions of “largeness” in analysis. A quite popular one is that of *prevalence*, which is discussed at length in [34]. In a Polish abelian group G , a set is prevalent if its complement A is Haar-null in the sense of Christensen [18], i.e. one can find a Borel probability measure ν on G such that $\nu(A+g) = 0$ for every $g \in G$. Some results concerning prevalence and hypercyclicity are proved in [10], and much more spectacular results regarding the size of the set of hypercyclic vectors are to be found in [31].

It is not incongruous to expect that if an operator T is ergodic in the Gaussian sense, then $HC(T)$ is *not* prevalent and even Haar-null. We are not able to prove this, but this is indeed true for a large class of ergodic weighted shifts, as shown by the following result.

Proposition 8.5. *Let X be a Banach space, and let $T \in \mathcal{L}(X)$. Assume that one can find $u \in X$ such that*

$$\sum_{n=0}^{\infty} \frac{1}{\|T^n(u)\|} < \infty.$$

Then $HC(T)$ is Haar-null.

Proof. Considering only the real-linear structure of X , we may assume that X is a real Banach space.

An efficient way of proving that a set $A \subset X$ is Haar-null is to exhibit some finite-dimensional subspace V of X such that

$$\forall x \in X \ \forall^{a.e.} v \in V : x + v \notin A,$$

where $\forall^{a.e.}$ refers to Lebesgue measure on V . In the terminology of [34], such a subspace V is called a *probe* for A .

We show that the one-dimensional subspace $V = \mathbb{R}u$ is a probe for $HC(T)$. So let $x \in X$ be arbitrary, set $\Lambda := \{\lambda \in \mathbb{R}; x + \lambda u \in HC(T)\}$, and let us prove that Λ has Lebesgue measure 0.

For any $\lambda \in \Lambda$, one can find arbitrary large $n \in \mathbb{N}$ such that

$$\|T^n(x) + \lambda T^n(u)\| \leq 1,$$

and hence such that

$$|\lambda - (\|T^n(x)\|/\|T^n(u)\|)| \leq \frac{1}{\|T^n(u)\|}.$$

Putting $a_n := \|T^n(x)\|/\|T^n(u)\|$, it follows that

$$|\lambda| \in \bigcap_{N \in \mathbb{N}} \bigcup_{n \geq N} \left[a_n - \frac{1}{\|T^n(u)\|}, a_n + \frac{1}{\|T^n(u)\|} \right]$$

for every $\lambda \in \Lambda$. In particular, the Lebesgue measure of Λ is not greater than

$$\inf_{N \in \mathbb{N}} 2 \sum_{n \geq N} \frac{1}{\|T^n u\|},$$

which is a complicated way to write 0. \square

Corollary 8.6. *Let $B_{\mathbf{w}}$ be a weighted backward shift on $X_p = \ell^p(\mathbb{N})$, $1 \leq p < \infty$ or $X_\infty = c_0(\mathbb{N})$. Assume that the weight sequence $\mathbf{w} = (w_n)_{n \geq 1}$ satisfies*

$$\sum_{n=1}^{\infty} \frac{1}{|w_1 \cdots w_n|^{p/p+1}} < \infty.$$

Then $HC(B_{\mathbf{w}})$ is Haar-null. This holds for example if $\liminf_{n \rightarrow \infty} |w_n| > 1$.

Proof. Set $w_0 := 0$ and denote by $(e_k)_{k \geq 0}$ the canonical basis of X_p . If $p < \infty$, consider the vector

$$u := \sum_{k=0}^{\infty} \frac{1}{|w_0 \cdots w_k|^{1/p+1}} e_k.$$

If $p = \infty$, put

$$u := \sum_{k=0}^{\infty} \frac{A_k}{w_0 \cdots w_k} e_k,$$

where (A_k) is any sequence of positive numbers such that $A_k = o(w_0 \cdots w_k)$ and $\sum_0^\infty 1/A_k < \infty$. \square

Thus, we see that if B is the usual (unweighted) backward shift, then $HC(\lambda B)$ is Haar-null in X_p for any $p \in [1, \infty]$ if $|\lambda| > 1$.

8.8. Some questions. To conclude the paper, we list a few “natural” questions, most of which have already been raised.

- (1) Is Theorem 2.6 true without assuming that the family \mathbf{S} is c_0 -like?
- (2) Is Theorem 2.8 true for $L^1(m)$ -mixing, or in the mild mixing case?
- (3) Let T be a hypercyclic operator on X whose \mathbb{T} -eigenvectors span a dense subspace of X . Is T ergodic in the Gaussian sense? Is T frequently hypercyclic?
- (4) Is every frequently hypercyclic Hilbert space operator necessarily ergodic (perhaps not in the Gaussian sense)?
- (5) Let $T \in \mathcal{L}(X)$, and assume that for any set $D \subset \mathbb{T}$ with empty interior, the linear span of $\bigcup_{\lambda \in \mathbb{T} \setminus D} \ker(T - \lambda)$ is dense in X . Is it possible to find a countable family of continuous \mathbb{T} -eigenvector fields $(E_i)_{i \in I}$, where each E_i is defined on some nontrivial closed arc $\Lambda_i \subset \mathbb{T}$, such that $\text{span}(\bigcup_{i \in I} E_i(\Lambda_i))$ is dense in X ?
- (6) Let H be a Hilbert space, and let $T \in \mathcal{L}(H)$ be mixing with respect to some Borel measure on H with full support. Is T mixing in the Gaussian sense?

- (7) On which Banach spaces is it possible to find ergodic operators (in the Gaussian sense) with no eigenvalues? The space X should of course not have cotype 2, but this is not enough: for example, if X is hereditarily indecomposable then there are no frequently hypercyclic operators at all on it, and hence no ergodic operators either ([55]). In fact, the question splits into two separate problems: (i) on which Banach spaces is it possible to find ergodic operators? (ii) what can be said if the space does not have cotype 2? We refer to [22] for general results regarding the first problem.
- (8) Is there an ergodic weighted shift on $c_0(\mathbb{N})$ with no unimodular eigenvalues?
- (9) Let $T \in \mathcal{L}(X)$ be ergodic in the Gaussian sense. Are the ergodic measures with full support dense in the set of all T -invariant measures (endowed with the usual Prokhorov topology)? See [56] for a positive answer in a completely different situation, and [19] for more in that direction.
- (10) Is it true that if T is an ergodic operator, then $HC(T)$ is Haar-null? Is this true at least for weighted shifts?

REFERENCES

- [1] J. Aaronson, *An introduction to infinite ergodic theory*. Mathematical Surveys and Monographs **50**, Amer. Math. Soc. (1997).
- [2] F. Albiac and N. Kalton, *Topics in Banach space theory*. Graduate Texts in Mathematics **233**, Springer (2006).
- [3] C. Badea and S. Grivaux, *Unimodular eigenvalues, uniformly distributed sequences and linear dynamics*. Adv. Math. **211** (2007), 766–793.
- [4] C. Badea and S. Grivaux, *Size of the peripheral point spectrum under power or resolvent growth conditions*. J. Funct. Anal. **247** (2007), 302–329.
- [5] C. Badea and V. Müller, *On weak orbits of operators*. Topology Appl. **156** (2009), 1381–1385.
- [6] F. Bayart and S. Grivaux, *Hypercyclicity and unimodular point spectrum*. J. Funct. Anal. **226** (2005), 281–300.
- [7] F. Bayart and S. Grivaux, *Frequently hypercyclic operators*. Trans. Amer. Math. Soc. **358** (2006), no. 11, 5083–5117.
- [8] F. Bayart and S. Grivaux, *Invariant Gaussian measures for operators on Banach spaces and linear dynamics*. Proc. Lond. Math. Soc. **94** (2007), 181–210.
- [9] F. Bayart and É. Matheron, *Dynamics of linear operators*. Cambridge Tracts in Mathematics **179**, Cambridge University Press (2009).
- [10] F. Bayart, É. Matheron and P. Moreau, *Small sets and hypercyclic vectors*. Comment. Math. Univ. Carolinae **49** (2008), 53–65.
- [11] F. Bayart and I. Z. Rusza, *Difference sets and frequently hypercyclic weighted shifts*. Ergodic Theory Dynam. Systems (to appear), available at <http://arxiv.org/abs/1305.2325>.
- [12] T. Bermúdez, A. Bonilla, J. Conejero and A. Peris, *Hypercyclic, topologically mixing and chaotic semigroups on Banach spaces*. Studia Math. **170** (2005), 57–75.
- [13] V. I. Bogachev *Gaussian measures*. Mathematical Surveys and Monographs **62**, Amer. Math. Soc. (1998).
- [14] A. Bonilla and K. -G. Grosse-Erdmann, *On a theorem of Godefroy and Shapiro*. Integral Equations Operator Theory **56** (2006), 151–162.
- [15] A. Bonilla and K. -G. Grosse-Erdmann, *Frequently hypercyclic operators and vectors*. Ergodic Theory Dynam. Systems **27** (2007), 383–404.
- [16] P. Brunovský and J. Komornik, *Ergodicity and exactness of the shift on $C[0, \infty]$ and the semiflow of a first order partial differential equation*. J. Math. Anal. Appl. **104** (1984), no. 1, 235–245.

- [17] S. A. Chobanyan, V. I. Tarieladze and N. N. Vakhania, *Probability distributions on Banach spaces*. Mathematics and its Applications (Soviet Series) **14**, Springer (1987).
- [18] J. P. R. Christensen, *On sets of Haar measure zero in Polish abelian groups*. Israel J. Math. **13** (1972), 255–260
- [19] Y. Coudène and B. Schapira, *Generic measures for hyperbolic flows on non-compact spaces*. Israel J. Math. **179** (2010), 157–172.
- [20] M. J. Cowen and R. G. Douglas, *Complex Geometry and Operator Theory*. Acta Math. **141** (1978), 187–261.
- [21] G. Da Prato, *An introduction to infinite-dimensional analysis*. Universitext, Springer (2006).
- [22] M. De la Rosa, L. Frerick, S. Grivaux and Alfredo Peris, *Frequent hypercyclicity, chaos, and unconditional Schauder decompositions*. Israel J. Math. **190** (2012), 389–399.
- [23] G. Debs and J. Saint Raymond, *Ensembles boréliens d’unicité et d’unicité au sens large*. Ann. Inst. Fourier, Grenoble **37** (1987), no. 3, 217–239.
- [24] W. Desch, W. Schappacher and G. F. Webb, *Hypercyclic and chaotic semigroups of linear operators*. Ergodic Theory Dynam. Systems **17** (1997), 793–819.
- [25] V. Devinck, *Strongly mixing operators on Hilbert spaces and speed of mixing*. Proc. London Math. Soc. **106** (2013), 1394–1434.
- [26] T. Eisner and S. Grivaux, *Hilbertian Jamison sequences and rigid dynamical systems*. J. Funct. Anal. **261** (2011), 2013–2052.
- [27] E. Flytzanis, *Unimodular eigenvalues and linear chaos in Hilbert spaces*. Geom. Funct. Anal. **5** (1995), 1–13.
- [28] E. Glasner, *Ergodic theory via joinings*. Mathematical Surveys and Monographs **101**, Amer. Math. Soc. (2003).
- [29] G. Godefroy and J. Shapiro, *Operators with dense, invariant, cyclic vector manifolds*. J. Funct. Anal. **98** (1991), 229–269.
- [30] S. Grivaux, *A new class of frequently hypercyclic operators*. Indiana Univ. Math. J. **60** (2011), 1177–1201.
- [31] S. Grivaux and M. Roginskaya, *On Read’s type operators on Hilbert spaces*. Int. Math. Res. Not. IMRN (2008), Art. ID rnm083, 42 pp.
- [32] K. -G Grosse-Erdmann, *Dynamics of linear operators*. Topics in operator theory and complex analysis 41–84, University of Malaga (2007).
- [33] K. -G Grosse-Erdmann and A. Peris, *Linear chaos*. Universitext, Springer (2011).
- [34] B. R. Hunt, T. Sauer and J. A. Yorke, *Prevalence: a translation invariant “almost every” on infinite-dimensional spaces*. Bull. Amer. Math. Soc. **27** (1992), no. 2, 217–238.
- [35] J. P. Kahane and R. Salem, *Ensembles parfaits et séries trigonométriques*. Hermann (1963).
- [36] S. Kahane, *On the complexity of sums of Dirichlet measures*. Ann. Inst. Fourier, Grenoble **43** (1993), no. 1, 111–123.
- [37] G. Kalisch, *On operators on separable Banach spaces with arbitrary prescribed point spectrum*. Proc. Amer. Math. Soc. **34** (1972) 207–208.
- [38] Y. Katznelson, *An introduction to harmonic analysis*. Dover (1976).
- [39] A. S. Kechris, *Classical descriptive set theory*. Graduate Texts in Mathematics **156**, Springer (1995).
- [40] A. S. Kechris and A. Louveau, *Descriptive set theory and the structure of sets of uniqueness*. London Math. Soc. Lecture Notes Series **128**, Cambridge University Press (1987).
- [41] A. Lasota, *Invariant measures and a linear model of turbulence*. Rend. Sem. Mat. Univ. Padova **61** (1979), 39–48.
- [42] R. Lyons, *Fourier–Stieltjes coefficients and asymptotic distribution modulo 1*. Ann. Math. **122** (1985) 155–170.
- [43] É. Matheron and M. Zelený, *Rudin-like sets and hereditary families of compact sets*. Fund. Math. **185** (2005), 97–116.
- [44] P. Mörters and Y. Peres, *Brownian motion*. Cambridge University Press (2010).
- [45] M. Murillo-Arcilla and A. Peris, *Strong mixing measures for linear operators and frequent hypercyclicity*. J. Math. Anal. Appl. **398** (2013), 462–465.

- [46] M. Murillo-Arcilla and A. Peris, *Strong mixing measures for C_0 -semigroups*. Preprint (2013), available at <http://arxiv.org/abs/1303.0327>
- [47] M. Nikula, *Frequent hypercyclicity of random entire functions for the differentiation operator*. Preprint (2012), available at <http://arxiv.org/abs/1209.6209>.
- [48] J. C. Oxtoby and S. M. Ulam, *On the existence of a measure invariant under a transformation*. Ann. Math. **40** (1939), 560–566.
- [49] J. C. Oxtoby and S. M. Ulam, *Measure-preserving homeomorphisms and metrical transitivity*. Ann. Math. **42** (1941), 874–920.
- [50] K. Petersen, *Ergodic theory*. Cambridge Studies in Advanced Mathematics **2**, Cambridge University Press (1983).
- [51] W. Rudin, *Functional analysis*. McGraw-Hill (1973).
- [52] R. Rudnicki, *Gaussian measure-preserving linear transformations*. Univ. Iagel Acta Math. **30** (1993), 105–112.
- [53] R. Rudnicki, *Chaos for some infinite-dimensional dynamical systems*. Math. Methods Appl. Sci. **27** (2004), no. 6, 723–738.
- [54] H. Salas, *Hypercyclic weighted shifts*. Trans. Amer. Math. Soc. **347** (1995), no. 3, 993–1004.
- [55] S. Shkarin, *On the spectrum of frequently hypercyclic operators*. Proc. Amer. Math. Soc. **137** (2009) no. 1, 123–134.
- [56] K. Sigmund, *Generic properties of invariant measures for axiom A diffeomorphisms*. Inventiones Math. **11** (1970), 99–109.
- [57] K. Zhu, *Operators in the Cowen-Douglas class*. Illinois J. Math. **44** (2000), 767–783.

CLERMONT UNIVERSITÉ, UNIVERSITÉ BLAISE PASCAL, LABORATOIRE DE MATHÉMATIQUES, BP 10448, F-63000 CLERMONT-FERRAND - CNRS, UMR 6620, LABORATOIRE DE MATHÉMATIQUES, F-63177 AUBIÈRE.

E-mail address: `Frederic.Bayart@math.univ-bpclermont.fr`

LABORATOIRE DE MATHÉMATIQUES DE LENS, UNIVERSITÉ D'ARTOIS, RUE JEAN SOUVRAZ S. P. 18, 62307 LENS.

E-mail address: `etienne.matheron@euler.univ-artois.fr`